

Ehresmann doubles and Drinfel'd doubles for Lie algebroids and Lie bialgebroids *

K. C. H. Mackenzie
Department of Pure Mathematics
University of Sheffield
Sheffield, S3 7RH
United Kingdom
K.Mackenzie@sheffield.ac.uk

December 18, 2006

Abstract

The word ‘double’ was used by Ehresmann to mean ‘an object X in the category of all X ’. Double categories, double groupoids and double vector bundles are instances, but the notion of Lie algebroid cannot readily be doubled in the Ehresmann sense, since a Lie algebroid bracket cannot be defined diagrammatically. In this paper we use the duality of double vector bundles to define a notion of double Lie algebroid, and we show that this abstracts the infinitesimal structure (at second order) of a double Lie groupoid.

We further show that the cotangent of either Lie algebroid in a Lie bialgebroid has a double Lie algebroid structure, and that a pair of Lie algebroid structures on dual vector bundles forms a Lie bialgebroid if and only if the structures which they canonically induce on their cotangents form a double Lie algebroid. In particular, the Drinfel'd double of a Lie bialgebra has a double Lie algebroid structure.

We also show that matched pairs of Lie algebroids, as used by J.–H. Lu in the classification of Poisson group actions, are in bijective correspondence with vacant double Lie algebroids.

*2000 *Mathematics Subject Classification*. Primary 53D17. Secondary 17B62, 17B66, 18D05, 22A22, 58H05.

INTRODUCTION

The word ‘double’ has two distinct usages. In [4] Drinfel’d introduced the classical double of a Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$, which combines $\mathfrak{g}$ and $\mathfrak{g}^*$ into a single Lie algebra $\mathfrak{g} \bowtie \mathfrak{g}^*$ which encodes fully the relations between $\mathfrak{g}$ and $\mathfrak{g}^*$. (We do not consider quantum doubles in this paper.)

In categorical language the word ‘double’ describes an object of type X in the category of all objects X . Thus a ‘double vector bundle’ is a vector bundle $D \rightarrow A$ in which both the total space and base space are objects in the category of vector bundles, say $D \rightarrow B$ and $A \rightarrow M$; the maps defining the structure of $D \rightarrow A$ must be morphisms in the category of vector bundles. (More detail is given below and in §1.) The notion of a double category, the fundamental prototype of this concept of double, was introduced and developed by Ehresmann [6]. Double categories should be distinguished from the more recent notions of 2-category and bicategory.

We show in this paper that the Drinfel’d double of a Lie bialgebra may be regarded as a double in the Ehresmann sense. Further, we show that the cotangent of (either bundle in) a Lie bialgebroid is a double in the Ehresmann sense, and that this construction reduces to the Drinfel’d double when the Lie bialgebroid is a Lie bialgebra.

The crucial step is to define a concept of double in the Ehresmann sense for Lie algebroids. The definition cannot be precisely of the type described above since the bracket of a Lie algebroid is not a map in the category of vector bundles. We use an indirect approach (Definition 3.1) and show, in §6, that this is an infinitesimal form of doubles of Ehresmann type.

A double Lie algebroid is first of all a double vector bundle as in Figure 1(a); that is, D




Figure 1.

has two vector bundle structures, on bases A and B , each of which is itself a vector bundle on base M , such that for each structure on D , the structure maps (projection, addition, scalar multiplication) are vector bundle morphisms with respect to the other structure, and the combination of the two projections, $D \rightarrow A \times_M B$, is a surjective submersion (see §1 for more detail). Two examples to keep in mind are the tangent prolongation of an ordinary vector bundle as in Figure 1(b) (see [1] for a classical treatment), and the cotangent double vector bundle as in Figure 1(c) (see [26]).

Now suppose that all four sides of Figure 1(a) have Lie algebroid structures. The problem is to define compatibility between the bracket structures on D , bearing in mind that the brackets are on different modules of sections. The key is the duality for double vector bundles described in §1. Although our compatibility condition cannot (as far as we know) be interpreted as a condition of the form ‘either bracket structure on (sections of) D is a morphism with respect

to the other', we show in Theorem 6.2 that this concept of double Lie algebroid includes the infinitesimal invariants of double Lie groupoids, which *are* defined by such conditions. We therefore feel justified in referring to it as a double of Ehresmannian type, or a categorical double.

In §4 we define a notion of double for Lie bialgebroids. A Lie bialgebroid (A, A^*) on base M [26] consists of Lie algebroid structures on a vector bundle and its dual, subject to a compatibility condition which can be most succinctly expressed by saying that the differential of each Lie algebroid acts as a derivation on the Schouten (or Gerstenhaber) algebra associated with the other [9].

Given a Lie bialgebroid (A, A^*) , the Lie algebroid structure on A induces a Poisson structure on A^* by an extension [3] of the duality between Lie algebras and linear Poisson structures on vector spaces. This Poisson structure in turn induces a Lie algebroid structure on $T^*A^* \rightarrow A^*$, defined in terms of the Poisson bracket of 1-forms. Likewise, the Lie algebroid structure on A^* induces a Lie algebroid structure on $T^*A \rightarrow A$. Using the canonical diffeomorphism $T^*A^* \rightarrow T^*A$ [26] we now have Lie algebroid structures on each of the four sides of Figure 1(c). This is a double Lie algebroid as defined in §3, the *cotangent double of (A, A^*)* .

Suppose now that we have Lie algebroid structures on A and A^* with no further assumption. As above, the Poisson structures on A^* and A induce Lie algebroid structures on $T^*A^* \rightarrow A^*$ and $T^*A \rightarrow A$ and we prove in §4 that if these constitute a double Lie algebroid structure on T^*A , then (A, A^*) is a Lie bialgebroid.

When M is a point, so that $A = \mathfrak{g}$ and A^* are Lie algebras, the cotangent manifold $T^*\mathfrak{g}$ is $\mathfrak{g} \times \mathfrak{g}^*$. The Lie algebroid structure on $\mathfrak{g} \times \mathfrak{g}^* \rightarrow \mathfrak{g}$ is the action, or transformation, Lie algebroid defined by the coadjoint action of $\mathfrak{g}^*$ on $\mathfrak{g}$; likewise the coadjoint action of $\mathfrak{g}$ on $\mathfrak{g}^*$ defines the Lie algebroid structure on $\mathfrak{g} \times \mathfrak{g}^* \rightarrow \mathfrak{g}^*$. These structures have not been much used, since $\mathfrak{g} \times \mathfrak{g}^*$ has the classical Drinfel'd double structure $\mathfrak{g} \bowtie \mathfrak{g}^*$, as a Lie algebra.

For an arbitrary Lie bialgebroid (A, A^*) there is no known Lie algebroid structure on $T^*A \rightarrow M$ (and indeed there is no natural vector bundle structure) which reduces to $\mathfrak{g} \bowtie \mathfrak{g}^*$ in the bialgebra case. Consequently there is no direct analogue for T^*A of the Manin triple theorem (as in [14]) which, for a pair of Lie algebra structures on dual vector spaces, gives a criterion for them to form a Lie bialgebra. Nonetheless Theorem 4.1 provides a criterion for a pair of Lie algebroid structures on dual vector bundles to form a Lie bialgebroid.

A Manin triple theorem for Lie bialgebroids which directly generalizes the Lie bialgebra result, was given by Liu, Weinstein, and Xu [11], whose notion of double is a Courant algebroid structure on the direct sum $A \oplus A^*$. Some comments on the differences between the two approaches are given at the end of §4.

The lack of a structure on $T^*A \rightarrow M$ is an inherent feature of the cotangent. A cotangent double vector bundle T^*A is locally isomorphic to $A \times_M A^* \times_M T^*M$ and is of the form $A \times_M A^*$ if and only if M is 0-dimensional.

We call a double Lie algebroid $(D; A, B; M)$ *vacant* if D is isomorphic to $A \times_M B$ as a double vector bundle. We prove in §5 that in this case there are induced actions ρ of A on B and σ of B on A , and these satisfy the equations

$$\begin{aligned} \rho_X([Y_1, Y_2]) &= [\rho_X(Y_1), Y_2] + [Y_1, \rho_X(Y_2)] + \rho_{\sigma_{Y_2}(X)}(Y_1) - \rho_{\sigma_{Y_1}(X)}(Y_2), \\ \sigma_Y([X_1, X_2]) &= [\sigma_Y(X_1), X_2] + [X_1, \sigma_Y(X_2)] + \sigma_{\rho_{X_2}(Y)}(X_1) - \sigma_{\rho_{X_1}(Y)}(X_2), \\ a(\sigma_Y(X)) - b(\rho_X(Y)) &= [b(Y), a(X)], \end{aligned} \tag{1}$$

for $X_1, X_2, X \in \Gamma A$, $Y_1, Y_2, Y \in \Gamma B$, where a and b are the anchors. Conversely, given such a pair of actions, $A \times_M B$ has a double Lie algebroid structure.

A pair of actions satisfying (1) is called a *matched pair* structure for A and B [31]. This notion of matched pair extends the concept introduced into geometry by Kosmann–Schwarzbach and Magri [10] under the name *extension bicroisée* or *twilled extension*, by Lu and Weinstein [14] as a *double Lie algebra*, and by Majid [28], with the term *matched pair* which we use. (In fact, forms of the concept had been found much earlier; see [37], [2] for references.) As in the case of Lie algebras, given a matched pair structure for Lie algebroids A and B , the direct sum bundle $A \oplus B$ has a Lie algebroid structure $A \bowtie B$ for which $A \oplus 0$ and $0 \oplus B$ are Lie subalgebroids, and the converse is also true [31].

We note here that Lu [12] showed that any Poisson action of a Poisson Lie group G on a Poisson manifold P gives rise to a matched pair of Lie algebroids, and the corresponding diagonal structure $T^*P \bowtie (P \times \mathfrak{g}^*)$ carries Drinfel'd's classification of Poisson homogeneous spaces [5]. In a future paper we will show that the corresponding construction for an action of a general Poisson groupoid gives rise to a double Lie algebroid which is not vacant, and so does not correspond to a matched pair.

* * * * *

There are two further points to be made about the concept of double Lie algebroid. Firstly, a natural global analogue is provided by the concept of double Lie groupoid [16]: we show in §6 that applying a second-order version of the Lie functor to a double Lie groupoid yields a double Lie algebroid. This subsumes various known differentiation results and raises a general integrability problem.

It is well known that Lie bialgebroids are the infinitesimal form of Poisson groupoids [36], [27]. In the case of Lie bialgebras a further step was achieved by Lu and Weinstein [13] who proved the existence of a symplectic double Lie groupoid integrating a Poisson Lie group. The double Lie algebroid of this symplectic double Lie groupoid is the cotangent double Lie algebroid described above.

For matched pairs of Lie algebras the global concept of matched pair of Lie groups was introduced into geometry at the same time [28], [14]. The global concept of a matched pair of Lie groupoids was given in [16]. In the Lie algebra case, various integrability results were given in the same papers, but little is known about the integrability of matched pairs of Lie algebroids [24]. As [13] shows, a matched pair of Lie algebras may fail to integrate to a matched pair of Lie groups, but nonetheless integrate to a double Lie groupoid.

The second point to be made is that the concept of double Lie algebroid offers, at least in some situations, a means of bypassing the problems caused by the lack of pullbacks in the Poisson category. It is well-known that the action groupoid arising from the action of a Poisson Lie group G on a Poisson manifold P is not a Poisson groupoid. It was shown in [22] that the induced action of $T^*G \rightrightarrows \mathfrak{g}^*$ on the dual $\mathfrak{p}$: $T^*P \rightarrow \mathfrak{g}^*$ of the infinitesimal action, called the *pith* in [22], gives rise to a groupoid object in the category of Lie algebroids which differentiates to Lu's matched pair of Lie algebroids. In the group case one may work directly with the matched pair but, as mentioned above, for a Poisson action of a Poisson groupoid, the corresponding structure is a double Lie algebroid which does not reduce to a matched pair.

Since the mid 1990s several authors have worked to provide a characterization of the cotangent double of a Lie bialgebroid in terms of super geometry [34], [33]. This has now

been achieved by Ted Voronov [35] who further gives a formulation of the general concept of double Lie algebroid presented here in terms of super geometry. Independently, Mehta [30] has found a super formulation of the related notion of $\mathcal{LA}$ -groupoid.

On a speculative level, one may ask what relation, if any, exists between this theory and the weak multiple category theory that is used in some approaches to field theory. The differences between double Lie groupoid theory [16], [20] and the various notions of weak 2-category or bicategory, are very considerable and it may be that there is no simple relationship. The expectation that a link may exist is perhaps based on the fact that the notion of ordinary Lie groupoid includes both the frame groupoids of differential geometry and the fundamental groupoid of a manifold, and thus has both a differential geometric and a topological significance. However, this may be misleading as regards multiple structures: the homotopy properties of double Lie groupoids present considerable problems, and the notion of weak 2-category does not seem to readily accept a smooth structure. Perhaps these matters can be resolved, but in any case the results of this paper, particularly in the light of §6, appear to show that multiple Lie groupoid theory is well suited to finite-dimensional smooth differential geometry.

* * * *

An announcement of some of the main results of this paper was published in [17], and an early version was arXived in [21]. Amongst the many changes between [21] and the current paper, the most striking is perhaps the simplification of the definition 3.1 of a double Lie algebroid, following the observation of Ted Voronov that in his super formulation [35], the analogue of Condition II in [21] follows from Conditions I and III. This has enabled many arguments to be considerably shortened.

We do not include coordinate representations in this paper. The use of intrinsic formulations keeps clear the nature of each structure and makes the lifting of results and definitions from one level to another much easier. For example, most of the canonical isomorphisms used in the paper have been introduced for manifolds or vector bundles, and are then lifted to various multiple structures. The functorial nature of these processes is less clear in coordinate formulations. However, a coordinate-based approach is included in [35].

The material of this paper has developed over a considerable number of years and I have benefited greatly from many conversations with Yvette Kosmann-Schwarzbach, Alan Weinstein, and Ping Xu over this time. Conversations with Ted Voronov at the *XXVth Workshop on Geometry and Mathematical Physics* at Białowieża in 2006 provided a crucial stimulus for this revision and I am very grateful to him, and to the organizers of the Białowieża workshop for providing, inter alia, the opportunity for extended conversations. I also wish to thank the organizers of the programme on *Gerbes, Groupoids and Quantum Field Theory* at the ESI in 2006, where important parts of this revision were written.

§ 1 DOUBLE VECTOR BUNDLES AND THEIR DUALITY

We begin by recalling the duality of double vector bundles which was introduced in [18], [8] using the duality which Pradines [32] had introduced for vector bundle objects in the category of Lie groupoids. See [15] for a fuller account.

The definition of *double vector bundle* given in the Introduction (see Figure 1(a)) is complete but needs elaboration. We denote the projections in Figure 1(a) by $q_A^D: D \rightarrow A$ and $q_B^D: D \rightarrow B$; these are vector bundle morphisms over the projections $q_B: B \rightarrow M$ and $q_A: A \rightarrow M$ respectively. If $d \in D$ has $q_A^D(d) = a$ and $q_B^D(d) = b$, and $q_A(a) = m$, then we say that d has *outline* $(d; a, b; m)$.

Consider four elements $d_1, \dots, d_4$ of D with projections $a_1, \dots, a_4$ in A and $b_1, \dots, b_4$ in B such that $a_1 = a_2$, $a_3 = a_4$, $b_1 = b_3$ and $b_2 = b_4$. Then the condition that each bundle projection from D is a morphism of vector bundles with respect to the other structure guarantees that both sides of

$$(d_1 \overset{+}{\underset{A}{+}} d_2) \overset{+}{\underset{B}{+}} (d_3 \overset{+}{\underset{A}{+}} d_4) = (d_1 \overset{+}{\underset{B}{+}} d_3) \overset{+}{\underset{A}{+}} (d_2 \overset{+}{\underset{B}{+}} d_4) \quad (2)$$

are defined. Here $\overset{+}{\underset{A}{+}}$ denotes the addition in $D \rightarrow A$ and $\overset{+}{\underset{B}{+}}$ the addition in $D \rightarrow B$. The condition that each addition be a vector bundle morphism with respect to the other structure then implies equality in (2). There are corresponding equations involving scalar multiplication. Equation (2) is the *interchange law for addition*.

Given $a \in A$ we denote the zero of $D \rightarrow A$ above a by $\tilde{0}_a$. Similarly the zero of $D \rightarrow B$ above $b \in B$ is denoted $\tilde{0}_b$. The zeros in $A \rightarrow M$ and $B \rightarrow M$ are denoted 0_m^A and 0_m^B . The interchange law implies that the zero $\tilde{0}_{0_m^A}$ for $D \rightarrow A$ and the zero $\tilde{0}_{0_m^B}$ for $D \rightarrow B$ are equal; this is denoted $\odot_m$.

The intersection of the kernels of the two projections defined on D is called the *core* of the double vector bundle D and denoted C . The vector bundle structures on D induce a common vector bundle structure on C with base M . The kernel of $q_B^D: D \rightarrow B$ is the pullback $q_A^!C$ of this vector bundle C across q_A , and the kernel of $q_A^D: D \rightarrow A$ is $q_B^!C$. In practice cores are often identified with structures which are not strictly subsets of D and in such cases we write $\bar{c}$ for the element of D which corresponds to a $c \in C$.

A *morphism of double vector bundles* from $(D; A, B; M)$ to $(D'; A', B'; M')$ is a quadruple of maps $\varphi: D \rightarrow D'$, $\varphi_A: A \rightarrow A'$, $\varphi_B: B \rightarrow B'$ and $\varphi_M: M \rightarrow M'$ such that (φ, φ_A) , (φ, φ_B) , (φ_A, φ_M) and (φ_B, φ_M) are morphisms of vector bundles. A morphism of double vector bundles induces a morphism of the cores $\varphi_C: C \rightarrow C'$ by restriction.

Regarding D as an ordinary vector bundle over A , we denote the dual bundle by $D \star A$. (This notation avoids problems with multiple subscripts later.) In addition to its standard structure on base A , this has a vector bundle structure on base C^* , the projection of which, denoted $\gamma_{C^*}^A$, is defined by

$$\langle \gamma_{C^*}^A(\Phi), c \rangle = \langle \Phi, \tilde{0}_a \overset{+}{\underset{B}{+}} \bar{c} \rangle \quad (3)$$

where $\Phi: (q_A^D)^{-1}(a) \rightarrow \mathbb{R}$, $a \in A_m$, and $c \in C_m$. The addition in $D \star A \rightarrow C^*$, which we denote by $\overset{+}{\underset{C^*}{+}}$, is defined by

$$\langle \Phi \overset{+}{\underset{C^*}{+}} \Phi', d \overset{+}{\underset{B}{+}} d' \rangle = \langle \Phi, d \rangle + \langle \Phi', d' \rangle. \quad (4)$$

The condition that Φ and Φ' project to the same element of C^* guarantees that this is well-defined. The zero of $D \star A \rightarrow C^*$ above $\kappa \in C_m^*$ is denoted $\tilde{0}_\kappa^{\star A}$ and is defined by

$$\langle \tilde{0}_\kappa^{\star A}, \tilde{0}_b +_A \bar{c} \rangle = \langle \kappa, c \rangle \quad (5)$$

where $c \in C_m$ and b is any element of B_m . The scalar multiplication is defined in a similar way. These two structures make $D \star A$ into a double vector bundle as in Figure 2(a), the




Figure 2.

vertical dual of D . The core of $D \star A$ identifies with B^* , with the core element $\bar{\psi}$ corresponding to $\psi \in B_m^*$ given by

$$\langle \bar{\psi}, \tilde{0}_b +_A \bar{c} \rangle = \langle \psi, b \rangle. \quad (6)$$

There is also a horizontal dual $D \star B$ with sides B and C^* and core A^* , as in Figure 2(b). There is now the following somewhat surprising result. A proof is in [15, Chap. 9].

Theorem 1.1 ([18, 3.1], [8, Thm. 16]) *There is a natural (up to sign) nondegenerate pairing between $D \star A \rightarrow C^*$ and $D \star B \rightarrow C^*$ given by*

$$|\Phi, \Psi| = \langle \Phi, d \rangle - \langle d, \Psi \rangle \quad (7)$$

where $\Phi \in D \star A$, $\Psi \in D \star B$ have $\gamma_{C^*}^A(\Phi) = \gamma_{C^*}^B(\Psi)$ and d is any element of D with $q_A^D(d) = \gamma_A^A(\Phi)$ and $q_B^D(d) = \gamma_B^B(\Psi)$.

The pairing on the LHS of (7) is over C^* , whereas the pairings on the RHS are over A and B respectively. The pairing induces isomorphisms of double vector bundles

$$Z_A: D \star A \rightarrow D \star B \star C^*, \quad \langle Z_A(\Phi), \Psi \rangle = |\Phi, \Psi|, \quad (8)$$

$$Z_B: D \star B \rightarrow D \star A \star C^*, \quad \langle Z_B(\Psi), \Phi \rangle = |\Phi, \Psi| \quad (9)$$

with $(Z_A) \star C^* = Z_B$. Note that Z_B induces the identity on both side bundles C^* and B and is $-\text{id}$ on the cores A^* , whereas Z_A induces the identity on the side bundles C^* and the cores B^* but is $-\text{id}$ on the side bundles A .

The example which follows is needed for §4.

Example 1.2 Given an ordinary vector bundle (A, q, M) there is the tangent double vector bundle of Figure 1(b). This example was used as early as the 1960s to provide an efficient formalism for connection theory in A ; see [1]. The core of TA consists of the vertical vectors along the zero section and identifies canonically with A . Dualizing the standard structure

$TA \rightarrow A$ gives the cotangent double vector bundle of Figure 1(c) with core T^*M . Dualizing the structure $TA \rightarrow TM$ gives a double vector bundle which we denote $(T^\bullet A; A^*, TM; M)$; this is canonically isomorphic to $(T(A^*); A^*, TM; M)$ under an isomorphism $I_A: T(A^*) \rightarrow T^\bullet A$ given by

$$\langle I_A(\mathcal{X}), \xi \rangle = \langle \mathcal{X}, \xi \rangle \quad (10)$$

where $\mathcal{X} \in T(A^*)$ and $\xi \in TA$ have $T(q_*)(\mathcal{X}) = T(q)(\xi)$, and $\langle \cdot, \cdot \rangle$ is the tangent pairing of $T(A^*)$ and TA over TM defined by

$$\langle \mathcal{X}, \xi \rangle = \left. \frac{d}{dt} \langle \varphi_t, a_t \rangle \right|_0,$$

for $\mathcal{X} = \left. \frac{d}{dt} \varphi_t \right|_0$ and $\xi = \left. \frac{d}{dt} a_t \right|_0$.

Applying 1.1 to $D = TA$ thus induces a pairing of T^*A and $T(A^*)$ over A^* given by

$$|\Phi, \mathcal{X}| = \langle \mathcal{X}, \xi \rangle - \langle \Phi, \xi \rangle$$

where $\Phi \in T^*A$ and $\mathcal{X} \in T(A^*)$ have $r(\Phi) = p_{A^*}(\mathcal{X})$, and $\xi \in TA$ is chosen so that $T(q)(\xi) = T(q_*)(\mathcal{X})$ and $p_A(\xi) = c_A(\Phi)$. (Here c_A is the projection of the standard cotangent bundle and $r: T^*A \rightarrow A^*$ is a particular case of (3).) This pairing is nondegenerate and so defines an isomorphism of double vector bundles $R: T^*A^* \rightarrow T^*A$ by the condition

$$|R(\mathcal{F}), \mathcal{X}| = \langle \mathcal{F}, \mathcal{X} \rangle$$

where the pairing on the RHS is the standard one of $T^*(A^*)$ and $T(A^*)$ over A^* . This R preserves the side bundles A and A^* but induces $-\text{id}: T^*M \rightarrow T^*M$ as the map of cores. It is an antisymplectomorphism with respect to the exact symplectic structures. In summary we now have the very useful equation

$$\langle \mathcal{F}, \mathcal{X} \rangle + \langle R(\mathcal{F}), \xi \rangle = \langle \mathcal{X}, \xi \rangle, \quad (11)$$

for $\mathcal{F} \in T^*A^*$, $\mathcal{X} \in T(A^*)$, $\xi \in TA$, where the pairings are over A^* , A and TM respectively. This equation was originally derived in [26, §5] by using the symplectic structures.

Now return to the general double vector bundle in Figure 1(a), denoting the core by C . Each vector bundle structure on D gives rise to a double cotangent bundle. These two double cotangents fit together into a triple structure as in Figure 3(a). This structure is a triple vector bundle in an obvious sense: each face is a double vector bundle and for each vector bundle structure on T^*D , the maps defining the structure are morphisms of double vector bundles. The left and rear faces of Figure 3(a) are the two cotangent doubles of the two structures on D and the top face may be regarded as the cotangent double of either of the two duals of D . (In all such triple diagrams, we take the oblique arrows to be coming out of the page.)

Figure 3(a) is, in a sense made precise in [23], the vertical dual of the tangent prolongation of D as given in Figure 3(b). Five of the six faces in 3(a) are double vector bundles of types considered already; it is only necessary to verify that the top face is a double vector bundle. The cores of the same five faces are known, and we take the core of the top face to be T^*C . Taking these cores in opposite pairs, together with a parallel edge, then gives three double vector bundles: the left and the right cores form $(T^*A; A, A^*; M)$, the back and the front cores form $(T^*B; B, B^*; M)$ and the up and the down cores form $(T^*C; C, C^*; M)$. Each of these *core double vector bundles* has core T^*M .




Figure 3.

§ 2 PRELIMINARY CASE

Consider a double vector bundle as in Figure 1(a). Before considering bracket structures, we need to consider two classes of sections of the bundle structures on D .

A section $\xi \in \Gamma_A D$ is a (*vertical*) *linear section* if there exists $X \in \Gamma B$ such that (ξ, X) is a vector bundle morphism from (A, q_A, M) to (D, q_B^D, B) .

Proposition 2.1 *If (ξ, X) is a linear section, then $\ell_\xi: D \star A \rightarrow \mathbb{R}$ defined by*

$$\Phi \mapsto \langle \Phi, \xi(\gamma_A^A(\Phi)) \rangle$$

is linear with respect to C^ as well as A , and the restriction of ℓ_ξ to the core of $D \star A$ is $\ell_X: B^* \rightarrow \mathbb{R}$.*

PROOF. Take elements $(\Phi_1; a_1, \kappa; m)$ and $(\Phi_2; a_2, \kappa; m)$ of $D \star A$. Their sum over C^* has outline $(\Phi_1 \overset{+}{\underset{C^*}{\Phi_2}}; a_1 + a_2, \kappa; m)$ and so, using first the linearity of ξ and then the definition of $\overset{+}{\underset{C^*}{\Phi_1}}$,

$$\begin{aligned} \ell_\xi(\Phi_1 \overset{+}{\underset{C^*}{\Phi_2}}) &= \langle \Phi_1 \overset{+}{\underset{C^*}{\Phi_2}}, \xi(a_1 + a_2) \rangle = \langle \Phi_1 \overset{+}{\underset{C^*}{\Phi_2}}, \xi(a_1) \overset{+}{\underset{B}{\xi(a_2)}} \rangle \\ &= \langle \Phi_1, \xi(a_1) \rangle + \langle \Phi_2, \xi(a_2) \rangle = \ell_\xi(\Phi_1) + \ell_\xi(\Phi_2). \end{aligned}$$

Scalar multiplication is similar. Given $\psi \in B_m^*$ the corresponding core element $\bar{\psi} \in D \star A$ is given by (6). Using this,

$$\ell_\xi(\bar{\psi}) = \langle \bar{\psi}, \xi(0_m^A) \rangle = \langle \psi, x(m) \rangle = \ell_x(\psi), \quad (12)$$

where $\ell_x: B^* \rightarrow \mathbb{R}$ denotes the linear map $\varphi \mapsto \langle \varphi, x(q_{B^*}(\varphi)) \rangle$, for any $x \in \Gamma B$. $\square$

Given a linear section (ξ, x) , denote by $\xi^\square$ the section of $D \star A \star C^* \rightarrow C^*$ which is defined by ℓ_ξ . Thus

$$\langle \xi^\square(\kappa), \Phi \rangle_{C^*} = \ell_\xi(\Phi) = \langle \Phi, \xi(a) \rangle_A \quad (13)$$

for $(\Phi; a, \kappa; m) \in D \star A$. It is easily seen that $(\xi^\square, x)$ is a linear section of $D \star A \star C^* \rightarrow C^*$.

There is thus a bijective correspondence between vertical linear sections of D and horizontal linear sections of $D \star A \star C^*$. In the case where D is the tangent of an ordinary vector bundle A , as in Figure 1(b), this is the correspondence between linear vector fields on A and linear vector fields on A^* (see [15, §3.4]).

Next, for any $c \in \Gamma C$, define a *core section* $\bar{c} \in \Gamma_A D$ by

$$\bar{c}(a) = \overline{c(m)} \underset{B}{+} \tilde{0}_a.$$

Proposition 2.2 *The vertical linear sections (ξ, X) and the core sections $\bar{c}$ for $c \in \Gamma C$, generate $\Gamma_A D$ as a $C^\infty(A)$ -module.*

PROOF. Firstly, given any $X \in \Gamma B$, there is a linear section (ξ, X) . To see this, let

$$S: D \rightarrow A \times_M B \times_M C$$

be a splitting of D ; that is, an isomorphism of double vector bundles to the double vector bundle for which the vertical structure is the pullback $q_A^!(B \oplus C)$ and the horizontal structure is the pullback $q_B^!(A \oplus C)$. Then the section

$$a \mapsto S^{-1}(a, X(m), 0_m),$$

where $m = q_A(a)$, is linear over X .

Now consider any section whatever $\xi \in \Gamma_A D$. Then by paracompactness $q_A^D \circ \xi: A \rightarrow B$ is a finite sum

$$q_A^D \circ \xi = \sum F_i \overline{X_i}$$

where $F_i \in C^\infty(A)$ and $\overline{X_i}$, for $X_i \in \Gamma B$, denotes the pullback section of $q_A^! B$. For each X_i choose a linear section (ξ_i, X_i) and write $\eta = \sum F_i \xi_i$.

Then $\xi - \eta$ projects to zero under q_A^D and is therefore $\bar{c}$ for some $c \in \Gamma C$. $\square$

Before addressing the general concept of double Lie algebroid, it will be helpful to deal with a very special case.

Definition 2.3 *An $\mathcal{LA}$ -vector bundle is a double vector bundle as in Figure 1(a) together with Lie algebroid structures on a pair of parallel sides, such that the structure maps of the other pair of vector bundle structures are Lie algebroid morphisms.*

For definiteness, take the Lie algebroid structures to be on $D \rightarrow A$ and $B \rightarrow M$. Write $\Delta_A: D \rightarrow TA$ and $b: B \rightarrow TM$ for the anchors.

Since $D \rightarrow A$ is a Lie algebroid, the dual $D \star A$ has a dual Poisson structure, which is linear with respect to the bundle structure over A .

Theorem 2.4 *The Poisson structure $\pi_{\star A}$ on $D \star A$ is also linear with respect to the bundle structure over C^* .*

PROOF. Since it is dual to a Lie algebroid structure, $\pi_{\star A}$ is characterized by the following equations.

$$\{\ell_\xi, \ell_\eta\} = \ell_{[\xi, \eta]}, \quad \{\ell_\xi, F \circ \gamma_A^A\} = \Delta_A(\xi)(F) \circ \gamma_A^A, \quad \{F_1 \circ \gamma_A^A, F_2 \circ \gamma_A^A\} = 0, \quad (14)$$

where ξ and η are arbitrary elements of $\Gamma_A D$ and $F_1, F_2 \in C^\infty(A)$. See [15, §10.3]. We need to prove corresponding equations with respect to the bundle structure over C^* . The functions $D \star A \rightarrow \mathbb{R}$ which are linear with respect to the bundle structure over C^* are determined by arbitrary sections of $D \star A \star C^* \rightarrow C^*$. As in Proposition 2.2, these are of two types.

Firstly, if $\mathcal{X}, \mathcal{Y} \in \Gamma_{C^*}(D \star A \star C^*)$ are linear sections, then $\mathcal{X} = \xi^\square$ and $\mathcal{Y} = \eta^\square$ for linear sections $(\xi, x), (\eta, y) \in \Gamma_A D$.

Lemma 2.5 *If (ξ, x) and (η, y) are linear sections of D , then $([\xi, \eta], [x, y])$ is also.*

PROOF. For a section ξ which projects under q_B^D to a section x , define a section $\xi \oplus \xi$ of $D \oplus_B D \rightarrow A \oplus_M A$ by $(\xi \oplus \xi)(a_1 \oplus a_2) = \xi(a_1) \oplus \xi(a_2)$. Then (ξ, x) is linear if and only if $\xi \oplus \xi$ projects to ξ under the horizontal addition $+_B: D \oplus_B D \rightarrow D$, which is a Lie algebroid morphism over $+: A \oplus_M A \rightarrow A$ by hypothesis. Since the two components of a $\xi \oplus \xi$ depend on each variable separately, we have $[\xi \oplus \xi, \eta \oplus \eta] = [\xi, \eta] \oplus [\xi, \eta]$ and the result follows. $\square$

So $\{\ell_{\mathcal{X}}^{C^*}, \ell_{\mathcal{Y}}^{C^*}\} = \{\ell_\xi, \ell_\eta\} = \ell_{[\xi, \eta]} = \ell_{[\xi, \eta]^\square}^{C^*}$ is also linear over C^* . Here we wrote $\ell_{\mathcal{X}}^{C^*}$ for clarity.

Secondly, any $\varphi \in \Gamma A^*$ induces a *core section* $\overline{\varphi}$ of $D \star A \star C^* \rightarrow C^*$ by

$$\overline{\varphi}(\kappa) = \widetilde{0}_\kappa^{*B} +_B \overline{\varphi(m)}, \quad \kappa \in C_m^*,$$

which in turn induces $\ell_{\overline{\varphi}}^{C^*}: D \star A \rightarrow \mathbb{R}$. A simple calculation shows that

$$\ell_{\overline{\varphi}}^{C^*} = \ell_\varphi \circ \gamma_A^A.$$

It now follows from (14) that

$$\{\ell_{\overline{\varphi}_1}^{C^*}, \ell_{\overline{\varphi}_2}^{C^*}\} = \{\ell_{\varphi_1} \circ \gamma_A^A, \ell_{\varphi_2} \circ \gamma_A^A\} = 0. \quad (15)$$

Next consider a linear section $\mathcal{X} = \xi^\square$ and $\varphi \in \Gamma A^*$. We have

$$\{\ell_{\mathcal{X}}^{C^*}, \ell_{\overline{\varphi}}^{C^*}\} = \{\ell_\xi, \ell_\varphi \circ \gamma_A^A\} = \Delta_A(\xi)(\ell_\varphi) \circ \gamma_A^A. \quad (16)$$

Since $\Delta_A(\xi)$ is a linear vector field on A , it maps the linear function ℓ_φ to a linear function.

To make this explicit, recall the notion of derivation on a vector bundle. A *derivation* on $E \rightarrow M$ is a linear differential operator Λ of order ≤ 1 for which there is a vector field X , depending on Λ , such that $\Lambda(f\mu) = f\Lambda(\mu) + X(f)\mu$ for all $f \in C^\infty(M)$ and all sections μ of E . For any vector bundle E there is a vector bundle $\mathcal{D}(E)$ the sections of which are the derivations, and with the anchor defined by $\Lambda \mapsto X$ and the usual bracket, $\mathcal{D}(E)$ is a Lie algebroid. See [15, §3.3] and references given there.

There is a bijective correspondence between derivations on E and linear vector fields on E^* , and in the present case we can define a derivation Λ_ξ on A^* such that

$$\Delta_A(\xi)(\ell_\varphi) = \ell_{\Lambda_\xi(\varphi)} \quad (17)$$

for all $\varphi \in \Gamma A^*$.

From 2.2 we know that $\ell_{\mathcal{X}}^{C^*}$, for $\mathcal{X}$ an arbitrary section of $D \star A \star C^* \rightarrow C^*$, is of the form

$$\sum F_i \ell_{\xi_i^\square}^{C^*} + \ell_{\overline{\varphi}}^{C^*}$$

where $F_i \in C^\infty(C^*)$. It follows from (15), (16), and the argument following Lemma 2.5, that the bracket of two such functions is another of the same type. This completes the proof that the bracket of any two functions $D \star A \star C^* \rightarrow \mathbb{R}$ which are linear over C^* , is also linear over C^* .

We next need to consider functions which are pullbacks across $\gamma_{C^*}^A: D \star A \rightarrow C^*$. Functions $C^* \rightarrow \mathbb{R}$ are of two types. Firstly any $c \in \Gamma C$ induces $\ell_c: C^* \rightarrow \mathbb{R}$. Secondly, any $f \in C^\infty(M)$ pulls back to $f \circ q_{C^*}: C^* \rightarrow \mathbb{R}$. It is easily seen that

$$\ell_c \circ \gamma_{C^*}^A = \ell_{\bar{c}}, \quad f \circ q_{C^*} \circ \gamma_{C^*}^A = f \circ q_A \circ \gamma_A^A. \quad (18)$$

For the bracket of two functions of the first type, we have $\{\ell_{\bar{c}_1}, \ell_{\bar{c}_2}\} = \ell_{[\bar{c}_1, \bar{c}_2]}$ from (14). Consider the horizontal scalar multiplication in D by some $t \neq 0, 1$. This defines an automorphism $D \rightarrow D$ over $A \rightarrow A$ and therefore induces a map of sections $t_B: \Gamma_A D \rightarrow \Gamma_A D$; the Lie algebroid condition then ensures that $[t_B(\xi), t_B(\eta)] = t_B([\xi, \eta])$ for all $\xi, \eta \in \Gamma_A D$. Now for $c \in \Gamma C$, $\bar{t}c = t_B(\bar{c})$ and so we have

$$\overline{[c_1, c_2]} = [t_B(\bar{c}_1), t_B(\bar{c}_2)] = [\bar{t}c_1, \bar{t}c_2] = \overline{t^2[c_1, c_2]}.$$

Thus the bracket of two functions $D \star A \rightarrow \mathbb{R}$ arising from sections of C must be zero.

That the bracket of two functions of the second type in (18) is zero follows immediately from (14).

Again from (14), we have

$$\{\ell_c \circ \gamma_{C^*}^A, f \circ q_{C^*} \circ \gamma_{C^*}^A\} = \{\ell_{\bar{c}}, f \circ q_{C^*} \circ \gamma_{C^*}^A\} = \Delta_A(\bar{c})(f \circ q_{C^*}) \circ \gamma_{C^*}^A.$$

The anchor $\Delta_A: D \rightarrow TA$ is a morphism of double vector bundles and therefore restricts to a map of the cores, which we denote by $\partial_A: C \rightarrow A$. Then Δ_A maps the core section $\bar{c}$ to the core section of TA corresponding to $\partial_A(c)$. The core sections of TA are the vertical vector fields on A and vertical vector fields map pullback functions to zero. (See [15, §3.4].) So $\{\ell_c \circ \gamma_{C^*}^A, f \circ q_{C^*} \circ \gamma_{C^*}^A\} = 0$.

It remains to show that brackets $\{\ell_{\mathcal{X}}^*, G \circ \gamma_{C^*}^A\}$, where $\mathcal{X}$ is a section of $D \star A \star C^* \rightarrow C^*$ and $G \in C^\infty(C^*)$, are of the form $G' \circ \gamma_{C^*}^A$. As before, it suffices to consider the cases where $\mathcal{X}$ is $\xi^\square$ or $\bar{\varphi}$ and G is ℓ_c or $f \circ q_{C^*}$.

Firstly, we have $\{\ell_{\ell^\square}^*, \ell_c \circ \gamma_{C^*}^A\} = \{\ell_\xi, \ell_{\bar{c}}\} = \ell_{[\xi, \bar{c}]}$. Now ξ projects to x under q_B^D and $\bar{c}$ projects to 0, so $[\xi, \bar{c}]$ also projects to 0. It is therefore $\bar{c}'$ for some $c' \in \Gamma C$ which we denote $Q_\xi(c)$; it is easily checked that $Q_\xi: \Gamma C \rightarrow \Gamma C$ is a derivation on C .

Secondly,

$$\{\ell_{\ell^\square}^*, f \circ q_{C^*} \circ \gamma_{C^*}^A\} = \{\ell_\xi, f \circ q_A \circ \gamma_A^A\} = \Delta_A(\xi)(f \circ q_A) \circ \gamma_A^A = x(f) \circ q_A \circ \gamma_A^A = x(f) \circ q_{C^*} \circ \gamma_{C^*}^A$$

which is of the required form.

In the third case we have

$$\{\ell_{\bar{\varphi}} \circ \gamma_A^A, \ell_{\bar{c}}\} = -\Delta_A(\bar{c})(\ell_{\bar{\varphi}}) \circ \gamma_A^A = -\partial_A(c)^\dagger(\ell_{\bar{\varphi}}) \circ \gamma_A^A,$$

where $\partial_A(c)^\dagger$ is the vertical vector field on A corresponding to the section $\partial_A(c)$. Since the flows of $\partial_A(c)^\dagger$ are by scalar multiplication, we have $\partial_A(c)^\dagger(\ell_{\bar{\varphi}}) = \langle \bar{\varphi}, \partial_A(c) \rangle \circ q_A$ and the result is of the required form.

Lastly, in the fourth case the bracket is zero, since both functions are pullbacks from A . This completes the proof of Theorem 2.4. $\square$

It follows that $D \star A \star C^* \rightarrow C^*$ has a Lie algebroid structure defined by

$$\ell_{[\mathcal{X}, \mathcal{Y}]}^{C^*} = \{\ell_{\mathcal{X}}^{C^*}, \ell_{\mathcal{Y}}^{C^*}\}, \quad e(\mathcal{X})(G) \circ \gamma_{C^*}^A = \{\ell_{\mathcal{X}}^{C^*}, G \circ \gamma_{C^*}^A\},$$

where e denotes the anchor, $\mathcal{X}, \mathcal{Y} \in \Gamma_{C^*}(D \star A \star C^*)$ and $G \in C^\infty(C^*)$.

From the proof of 2.4 we have, in particular, that

$$\begin{aligned} [\xi^\square, \eta^\square] &= [\xi, \eta]^\square, & [\xi^\square, \bar{\varphi}] &= \overline{\Lambda_\xi(\varphi)}, & [\bar{\varphi}, \bar{\psi}] &= 0, & e(\xi^\square)(\ell_c) &= \ell_{Q_\xi(c)}, \\ e(\xi^\square)(f \circ q_{C^*}) &= x(f) \circ q_{C^*}, & e(\bar{\varphi})(\ell_c) &= -\langle \varphi, \partial_A(c) \rangle \circ q_{C^*}, & e(\bar{\varphi})(f \circ q_{C^*}) &= 0, \end{aligned} \quad (19)$$

where (ξ, x) and (η, y) are linear sections of $D \rightarrow A$, $\varphi, \psi \in \Gamma A^*$, $c \in \Gamma C$ and $f \in C^\infty(M)$. The operators $\Lambda_\xi: \Gamma A^* \rightarrow \Gamma A^*$ and $Q_\xi: \Gamma C \rightarrow \Gamma C$ are defined in the course of the proof.

Notice that, since the relationship between D and $D \star A \star C^*$ is reciprocal, there is a similar set of equations for the bracket on $D \rightarrow A$.

For any vector bundle $V \rightarrow M$ with a linear Poisson structure π , the Poisson anchor $\pi^\#: T^*V \rightarrow TV$ is a morphism of double vector bundles over V and the anchor $a: V^* \rightarrow TM$ of the dual Lie algebroid structure. Taking the dual of $\pi^\#$ over V (that is, the usual dual) exchanges the side and core structures with the other's duals, so that $(\pi^\#)^*$ has core morphism a^* . Since $(\pi^\#)^* = -\pi^\#$ it follows that the core morphism of $\pi^\#$ is $-a^*$. See [15, 9.2.1].

This argument applies to each vector bundle structure on $D \star A$. Denote the Poisson anchor associated to $\pi_{\star A}$ by $\pi^{\#A}$. Figure 4 shows the cotangent and tangent triple vector bundles for $D \star A$, which are the domain and target for $\pi^{\#A}$.




Figure 4.

Since $\pi_{\star A}$ is linear over A , $\pi^{\#A}: T^*(D \star A) \rightarrow T(D \star A)$ is a morphism of the left hand faces in Figure 4, and the map $D \rightarrow TA$ is the anchor Δ_A .

Likewise, $\pi^{\#A}$ is a morphism of double vector bundles for the rear faces of Figure 4 and the corner map $D \star A \star C^* \rightarrow TC^*$ is the anchor e for the Lie algebroid structure on $D \star A \star C^*$.

It follows by a simple argument that $\pi^{\#A}$ is a morphism of triple vector bundles. The corner map $B \rightarrow TM$ is the anchor b and the other four corner maps are identities.

In effect, Theorem 2.4 shows that dualizing an $\mathcal{LA}$ -vector bundle along its Lie algebroid structure gives rise to a *Poisson double vector bundle*; that is, to a double vector bundle with

a Poisson structure for which the Poisson anchor is a morphism of triple vector bundles. The converse may also be proved.

§ 3 ABSTRACT DOUBLE LIE ALGEBROIDS

We now turn to the general notion of double Lie algebroid. Before giving the definition, we need some preliminaries.

Again consider a double vector bundle as in Figure 1(a). We now assume that there are Lie algebroid structures on all four sides and that each pair of parallel sides forms an $\mathcal{LA}$ -vector bundle.

Denote the four anchors by $\Delta_A: D \rightarrow TA$, $\Delta_B: D \rightarrow TB$, and $b: B \rightarrow TM$, $a: A \rightarrow TM$. As usual we denote all four brackets by $[\cdot, \cdot]$; the notation for elements will make clear which structure we are using.

The anchors thus give morphisms of double vector bundles

$$\begin{aligned} (\Delta_A; \text{id}, b; \text{id}): (D; A, B; M) &\rightarrow (TA; A, TM; M), \\ (\Delta_B; a, \text{id}; \text{id}): (D; A, B; M) &\rightarrow (TB; TM, B; M) \end{aligned}$$

and so define morphisms of their cores; denote these by $\partial_A: C \rightarrow A$ and $\partial_B: C \rightarrow B$.

From §2 we know that $D \star A \star C^*$ is a Lie algebroid over C^* . Likewise, the Lie algebroid structure on $D \rightarrow B$ induces a Poisson structure on $D \star B$ which is linear over C^* and therefore induces a Lie algebroid structure on $D \star B \star C^*$ with base C^* .

The bundles $D \star A \star C^*$ and $D \star B \star C^*$ are dual over C^* under the pairing:

$$[\mathcal{X}, \mathcal{Y}] = \langle \mathcal{X}, Z_A^{-1}(\mathcal{Y}) \rangle = \langle Z_B^{-1}(\mathcal{X}), \mathcal{Y} \rangle.$$

Definition 3.1 *A double Lie algebroid is a double vector bundle as in Figure 1(a), equipped with Lie algebroid structures on all four sides such that each pair of parallel sides forms an $\mathcal{LA}$ -vector bundle, and such that the Lie algebroid structures on $D \star A \star C^* \rightarrow C^*$ and $D \star B \star C^* \rightarrow C^*$ form a Lie bialgebroid.*

In practice we will use Z_A or Z_B to transfer the Lie algebroid structure from, for example, $D \star B \star C^*$ to $D \star A$.

We recall the notion of Lie bialgebroid, which was introduced in [26] as an abstraction of the infinitesimal structure of a Poisson groupoid [36]. The conventions used here follow [15].

Definition 3.2 *Let A be a Lie algebroid on base M and suppose that A^* is also equipped with a structure of Lie algebroid. These structures constitute a Lie bialgebroid if*

$$d_*[X, Y] = [d_*X, Y] + [X, d_*Y] \tag{20}$$

for all $X, Y \in \Gamma A$.

In this definition d_* is the coboundary operator for A^* and $[\cdot, \cdot]$ is the Schouten bracket of multisections of A . See [15], for example.

A Lie bialgebra is a Lie bialgebroid for which M is a single point. For a Poisson manifold M , the cotangent Lie algebroid structure on T^*M makes (T^*M, TM) a Lie bialgebroid. The following material, which is needed below, can be found in [26] and [15, Chapter 12].

Suppose that (A, A^*) is a Lie bialgebroid on base M and denote the anchors by a and a_* . Then we take the Poisson structure on M to be $\pi_M^\# = a_* \circ a^*$. It follows that a is a Poisson map (to the tangent lift Poisson structure on TM ; see below) and a_* is anti-Poisson.

The concept of Lie bialgebroid is self-dual: if (A, A^*) is a Lie bialgebroid, then (A^*, A) is also ([26, 3.10], [9], or see [15]). However (A^*, A) will induce on M the opposite of the Poisson structure induced by (A, A^*) .

In this section we need the following *morphism criterion* for a Lie bialgebroid ([26, 6.2], or see [15, §12.2]).

Theorem 3.3 *Let A be a Lie algebroid on M such that its dual vector bundle A^* also has a Lie algebroid structure. Denote their anchors by a and a_* . Then (A, A^*) is a Lie bialgebroid if and only if*

$$T^*(A^*) \xrightarrow{R} T^*(A) \xrightarrow{\pi_A^\#} TA \quad (21)$$

is a Lie algebroid morphism over a_ , where the domain $T^*(A^*) \rightarrow A^*$ is the cotangent Lie algebroid induced by the Poisson structure on A^* , and the target $TA \rightarrow TM$ is the tangent prolongation of A .*

The tangent prolongation Lie algebroid structure on $TA \rightarrow TM$ is best thought of in terms of the dual structure. For any Poisson manifold P there is a *tangent lift* Poisson structure on TP for which the anchor is given by

$$\pi_{TP}^\# = J_P \circ T(\pi_P^\#) \circ \Theta_P^{-1} \quad (22)$$

where $J_P: T^2P \rightarrow T^2P$ is the canonical involution for the manifold P and $\Theta_P: T(T^*P) \rightarrow T^*(TP)$ was named the *Tulczyjew diffeomorphism of P* in [15, p.395] (Θ was denoted J' in [26]). For any manifold P , either of the two formulas

$$\Theta_P = R_{TM} \circ (\delta\nu)^\flat = J_M^* \circ I_{TM} \quad (23)$$

may be taken as the definition of Θ_P . Here $\delta\nu$ is the canonical symplectic structure on T^*P .

If $P = A^*$ is the dual of a Lie algebroid then the tangent lift Poisson structure is linear with respect to the bundle $T(A^*) \rightarrow TM$ and therefore induces a Lie algebroid structure on $TA \rightarrow TM$ via the tangent pairing $\langle\langle \cdot, \cdot \rangle\rangle$. The anchor is $J_M \circ T(a)$. See [26, 5.6] or [15, 10.3.14, 9.7.1].

With these preliminaries established, consider a double Lie algebroid $(D; A, B; M)$. We use Z_A to transfer the Lie algebroid structure from $D \star B \star C^*$ to $D \star A$. From the morphism criterion, we know that

$$\pi^{A^\#} \circ R: T^*(D \star A \star C^*) \rightarrow T(D \star A)$$

is a morphism of Lie algebroids over $e: D \star A \star C^* \rightarrow TC^*$. As with any Lie bialgebroid, it is also a morphism of double vector bundles, with the other side morphism being the identity of $D \star A$ and the core morphism being $e^*: T^*C^* \rightarrow D \star A$.

From §2 we know that e is a morphism of double vector bundles, namely of the two right-hand faces in Figure 4, over C^* and $b: B \rightarrow TM$.

Lemma 3.4 *The core of $e: D \star A \star C^* \rightarrow TC^*$ is $-\partial_A^*: A^* \rightarrow C^*$.*

PROOF. The dual of $\pi^{\#A}$ over $D \star A$ is $-\pi^{\#A}$. In the dualization process, the two triple vector bundles in Figure 4 are interchanged, but the positioning of the sides and cores is permuted. In particular, the cores of the right-hand faces and of the front faces are exchanged. The map of the front cores induced by $\pi^{\#A}$ is ∂_A , by definition. $\square$

The process of dualizing a triple structure is covered in [23]. Lemma 3.4 can also be proved by using (19).

Dualizing e over C^* it follows that e^* is a morphism of double vector bundles over C^* and $-\partial_A$, with core morphism $b^*: T^*M \rightarrow B^*$.

The Poisson structure induced on C^* by the Lie bialgebroid $(D \star A \star C^*, D \star A)$ has associated anchor

$$\pi_{C^*}^{\#} = e_* \circ e^*: T^*C^* \rightarrow TC,$$

where $e_*: D \star A \rightarrow TC^*$ is the anchor of $D \star A \rightarrow C^*$. Denoting the anchor of $D \star B \star C^* \rightarrow C^*$ by f , we have $e_* = f \circ Z_A$. The arguments applied above to e apply equally to f and so $f: D \star B \star C^* \rightarrow TC^*$ is a morphism of double vector bundles over C^* and $-a$, with core morphism $-\partial_B^*$. Thus $\pi_{C^*}^{\#}$ is a morphism of double vector bundles and therefore induces on C the structure of a Lie algebroid, for which the anchor is the side morphism $(-a) \circ (-\partial_A)$.

The core morphism for $\pi_{C^*}^{\#}$ is $(-\partial_B^*) \circ b^*$. The skew-symmetry of a Poisson anchor implies that its core morphism is the negative dual of its non-identity side morphism, and so we have $a \circ \partial_A = b \circ \partial_B$.

Proposition 3.5 *The maps $\partial_A: C \rightarrow A$ and $\partial_B: C \rightarrow B$ are Lie algebroid morphisms.*

We need two preparatory lemmas. The first is from [15, 10.3.6] and is proved by the same method as the second.

Lemma 3.6 *For any Lie algebroid V on M , the bundle projection $T^*V^* \rightarrow V$ is a morphism of Lie algebroids over the bundle projection $V^* \rightarrow M$.*

Lemma 3.7 *The projections $D \star A \star C^* \rightarrow B$ and $D \star B \star C^* \rightarrow A$ are morphisms of Lie algebroids over the bundle projection $C^* \rightarrow M$.*

PROOF. Denote the projections by $Q: D \star A \star C^* \rightarrow C^*$ and $q: C^* \rightarrow M$. The anchor condition $b \circ Q = T(q) \circ f$ follows from the fact that the Poisson anchor for $D \star A$, namely $T^*(D \star A) \rightarrow T(D \star A)$, is not only a morphism of double vector bundles, but a morphism of triple vector bundles.

Since q is a surjective submersion and Q is fibrewise surjective, it is sufficient to check that if $\mathcal{X}$ projects to x and $\mathcal{Y}$ projects to y , then $[\mathcal{X}, \mathcal{Y}]$ projects to $[x, y]$. By 2.2, it is sufficient to consider the cases of linear sections and core sections.

For two linear sections (ξ, x) and (η, y) of $D \rightarrow A$ we have from the proof of Theorem 2.4 that $[\xi, \eta]$ is linear over $[x, y]$ and $[\xi^\square, \eta^\square] = [\xi, \eta]^\square$. Likewise, for $\varphi, \psi \in \Gamma A^*$, the core sections project to zero and $[\overline{\varphi}, \overline{\psi}] = 0$ does also.

Lastly, for a linear section (ξ, x) and $\varphi \in \Gamma A^*$, we have from (19) that $[\xi^\square, \overline{\varphi}]$ is the core section $\Lambda_\xi(\varphi)$, and this projects to zero.

The result for $D \star B \star C^* \rightarrow A$ is proved in the same way. $\square$

PROOF OF 3.5: As noted above, $e^*: T^*C^* \rightarrow D \star A$ is a morphism of double vector bundles over C^* and $-\partial_A: C \rightarrow A$. As for any Lie bialgebroid, e^* is also a morphism of Lie algebroids

over C^* . Now the projection $T^*C^* \rightarrow C$ is a morphism of Lie algebroids, and the projection $D \star A \rightarrow A$ is an anti-morphism of Lie algebroids (because Z_A is $-\text{id}$ on A). Both projections are surjective submersions and so determine the Lie algebroid structures on their targets. It follows that $-\partial_A$ is an anti-morphism of Lie algebroids. $\square$

The general process used here, of quotienting a known morphism of Lie algebroids, is based on the following result. A *fibration of Lie algebroids* is a morphism $\nu: A \rightarrow A'$ over $p: M \rightarrow M'$ such that p is a surjective submersion and ν is fibrewise surjective.

Proposition 3.8 *Consider Lie algebroids A , A' and B on bases M , M' and N , and let $\varphi: A \rightarrow A'$, $\psi: A' \rightarrow B$ and $\nu: A \rightarrow A'$ be morphisms of vector bundles over f, g and p , with $\varphi = \psi \circ \nu$.*

If φ is a morphism of Lie algebroids and ν is a fibration of Lie algebroids, then ψ is a morphism of Lie algebroids.

This follows from the general theory of quotients of Lie algebroids [15, §4.4], but is easy to prove directly.

Observe that Proposition 3.5 cannot be proved by mapping sections of C to core sections of $D \rightarrow A$. In the case of an $\mathcal{LA}$ -vector bundle there is a simple relation (equation (18)), but this does not hold for a general double Lie algebroid. This may be seen by considering the iterated tangent bundle $T^2M = T(TM)$ of any manifold; the bracket of two vertical lift vector fields (which are the core sections in this case) is always zero.

The next result, that the anchors Δ_A and Δ_B are morphisms of Lie algebroids with respect to both structures, was taken as part of the definition of a double Lie algebroid in [21]. I am grateful to Ted Voronov for having shown [35], using super methods, that it is a consequence of the other conditions in the definition. The proof which follows uses standard methods.

Theorem 3.9 *The anchor $\Delta_A: D \rightarrow TA$ is a Lie algebroid morphism over $b: B \rightarrow TM$, and the anchor $\Delta_B: D \rightarrow TB$ is a Lie algebroid morphism over $a: A \rightarrow TM$.*

$$\begin{array}{ccccc}
 T^*(D \star A \star C^*) & \xrightarrow{\pi^{\#A} \circ R} & T(D \star A) & & \\
 \downarrow & \searrow \nu & \downarrow \Delta_A & \searrow T(\nu') & \\
 D & \xrightarrow{\quad} & D & \xrightarrow{\quad} & TA \\
 \downarrow & \searrow e & \downarrow & \searrow & \downarrow \\
 D \star A \star C^* & \xrightarrow{\quad} & TC^* & \xrightarrow{\quad} & TM \\
 \downarrow & \searrow & \downarrow & \searrow & \downarrow \\
 B & \xrightarrow{\quad} & B & \xrightarrow{\quad} & TM
 \end{array}$$

Figure 5.

PROOF. Figure 5 shows part of the outline of the map $\pi^{\#A} \circ R: T^*(D \star A \star C^*) \rightarrow TC^*$. The dashed lines indicate that this should be read as a morphism of double structures, from left to right, and emphasize that it is not a triple structure,

Note first that every vertical arrow is the bundle projection of a Lie algebroid. Next, $\pi^{\#A} \circ R$ is a morphism of Lie algebroids over e , by the morphism criterion. Next, v and $T(\nu')$ are fibrations of Lie algebroids. That $\nu': D \star A \rightarrow A$ is a morphism over $q_*: C^* \rightarrow M$ follows from Lemma 3.7, and so its tangent prolongation is also. That ν is a fibration follows from 3.6 by using $D \star A \star C^* \cong D \star B$ and taking $V = D \rightarrow B$.

The result now follows by Proposition 3.8. $\square$

In the sections which follow we consider three classes of examples of double Lie algebroids.

§ 4 THE COTANGENT DOUBLE OF A LIE BIALGEBROID

The notion of Lie bialgebroid is recalled in the previous section.

Consider a Lie algebroid A on M together with a Lie algebroid structure on the dual, not *a priori* related to that on A . The structure on A^* induces a Poisson structure on A , and this gives rise to a cotangent Lie algebroid $T^*A \rightarrow A$. Equally, the Lie algebroid structure on A induces a Poisson structure on A^* and this gives rise to a cotangent Lie algebroid $T^*A^* \rightarrow A^*$. We transfer this latter structure to $T^*A \rightarrow A^*$ via R . There are now Lie algebroid structures on each of the four sides of the double vector bundle $D = T^*A$ of Figure 6(a).

Theorem 4.1 *Let A be a Lie algebroid on M such that its dual vector bundle A^* also has a Lie algebroid structure. Then (A, A^*) is a Lie bialgebroid if and only if $D = T^*A$, with the structures just described, is a double Lie algebroid.*

PROOF. First assume merely that A^* is a Lie algebroid. Consider Figure 1(c) with the corresponding Lie algebroid structures on the horizontal sides.

Lemma 4.2 *Figure 1(c) with the Lie algebroid structures just described, is an $\mathcal{LA}$ -vector bundle.*

PROOF. It must be proved that the vector bundle operations are Lie algebroid morphisms. That the bundle projection $T^*A \rightarrow A^*$ is a Lie algebroid morphism follows from Lemma 3.6. That the addition map is a Lie algebroid morphism follows from the fact that $\delta\ell_\varphi + \delta\ell_\psi = \delta\ell_{\varphi+\psi}$ for $\varphi, \psi \in \Gamma A^*$. The other conditions are similar. $\square$

Now assume that (A, A^*) is a Lie bialgebroid. The vertical structure on D is the cotangent Lie algebroid structure for the Poisson structure on A . Dualizing over A we have Figure 6(b) and the Poisson structure on TA is the tangent lift of the Poisson structure on A . By §2 this Poisson structure is also linear over TM . The dual $D \star A \star TM$ is shown in Figure 6(c). The map $I_A: T(A^*) \rightarrow T^*A$ described in Example 1.2 is an isomorphism of double vector bundles to $(T(A^*); A^*, TM; M)$ and by [15, 10.3.14] it is an isomorphism of Lie algebroids over TM to the tangent prolongation of $A^* \rightarrow M$.

Now consider the horizontal Lie algebroid structure in Figure 6(a). This was transported via R from the cotangent Lie algebroid of A^* . By the same argument as in the vertical case, the induced Lie algebroid structure on $D \star A^* \star TM \cong TA \rightarrow TM$ is isomorphic to the tangent prolongation Lie algebroid of $A \rightarrow M$.

We must now prove that :

Proposition 4.3 *For (A, A^*) a Lie bialgebroid on M , and the structures just described, the pair (TA, T^*A) is a Lie bialgebroid on TM .*




Figure 6.

PROOF. We use the morphism criterion 3.3. We must prove that

$$T^*(T^\bullet A) \xrightarrow{R_\bullet} T^*(TA) \xrightarrow{\pi_{TA}^\#} T^2 A \quad (24)$$

is a morphism of Lie algebroids over the anchor $J_M \circ T(a_*) \circ I_A^{-1}: T^\bullet A \rightarrow T^2 M$ of $T^\bullet A$.

Here $R_\bullet$ is the canonical map R for $TA \rightarrow TM$. The domain of (24) is the cotangent Lie algebroid for the Poisson structure on $T^\bullet A$. The target of (24) is the iterated tangent prolongation of the Lie algebroid structure of A .

We need two lemmas. The first is easily proved (see [15, 9.6.4]). The Θ is the Tulczyjew diffeomorphism defined in (23).

Lemma 4.4 *Let M be any manifold and write $p: TM \rightarrow M$ and $c: T^*M \rightarrow M$ for the bundle projections. Then for $\mathcal{X} \in T(T^*M)$ and $\xi \in T(TM)$ with $T(c)(\mathcal{X}) = T(p)(\xi)$,*

$$\langle\langle \mathcal{X}, \xi \rangle\rangle = \langle \Theta(\mathcal{X}), J(\xi) \rangle.$$

In the next lemma, the map $I_A^\times: T^*(T^\bullet A) \rightarrow T^*(TA^*)$ is the diffeomorphism of the cotangents induced by the diffeomorphism I_A .

Lemma 4.5 $R_\bullet = \Theta_A \circ T(R_A) \circ \Theta_{A^*}^{-1} \circ I_A^\times$.

PROOF. We begin by calculating $R_\bullet$ in terms of (11). Take $F \in T^*(T^\bullet A)$ and $\xi \in T^2 A$ projecting to the same point of TA . Then

$$\langle R_\bullet(F), \xi \rangle_{TA} = \langle\langle \mathcal{X}, \xi \rangle\rangle_{T^2 M} - \langle F, \mathcal{X} \rangle_{T^\bullet A}, \quad (25)$$

where $\mathcal{X} \in T(T^\bullet A)$ has the requisite projections. Write $\Phi = \Theta_{A^*}^{-1}(I_A^\times(F)) \in T(T^*A^*)$. Then

$$\langle R_\bullet((I_A^\times)^{-1}(\Theta_{A^*}(\Phi))), \xi \rangle_{TA} = \langle\langle \mathcal{X}, \xi \rangle\rangle_{T^2 M} - \langle (I_A^\times)^{-1}(\Theta_{A^*}(\Phi)), \mathcal{X} \rangle_{T^\bullet A},$$

Define $\Psi = T(I_A)^{-1}(\mathcal{X}) \in T^2(A^*)$. Then the first term on the RHS is

$$\langle\langle T(I_A)(\Psi), \xi \rangle\rangle_{T^2 M} = \frac{d}{dt} \langle I_A(\Psi_t), \xi_t \rangle_{TM} \Big|_0 = \frac{d}{dt} \langle\langle \Psi_t, \xi_t \rangle\rangle_{TM} \Big|_0.$$

where Ψ_t and ξ_t are curves in TA^* and TA with tangents Ψ and ξ . Denote this expression by $\langle\langle\langle \Psi, \xi \rangle\rangle\rangle$; this is a pairing of $T^2(A^*)$ and $T^2(A)$ over $T^2 M$.

The second term on the RHS of (25) is equal to $\langle \Theta_{A^*}(\Phi), T(I_A^{-1})(\mathcal{X}) \rangle_{T^*A}$. By Lemma 4.4, this is $\langle \langle \Phi, J_{A^*}(\Psi) \rangle \rangle_{TA^*}$.

Now consider $\Theta_A \circ T(R)$. Using 4.4 again,

$$\langle \Theta_A(T(R)(\Phi)), \xi \rangle_{TA} = \langle T(R)(\Phi), J_A(\xi) \rangle_{TA} = \left. \frac{d}{dt} \langle R(\Phi_t), \eta_t \rangle \right|_0,$$

where Φ_t is a curve in T^*A^* with tangent Φ and η_t is a curve in TA with tangent $J_A(\xi)$. Now

$$\langle R(\Phi_t), \eta_t \rangle = \langle W_t, \eta_t \rangle_{TM} - \langle \Phi_t, W_t \rangle_{A^*}$$

where $W_t \in TA^*$ is any element with the required outline. A careful checking of the outlines of the various elements shows that we can take W_t to be an integral curve of $J_{A^*}(\Psi)$. It is then clear that

$$\left. \frac{d}{dt} \langle \Phi_t, W_t \rangle_{A^*} \right|_0 = \langle \langle \Phi, J_{A^*}(\Psi) \rangle \rangle_{TA^*}.$$

The result now follows from the next lemma. $\square$

Lemma 4.6 *Let $A \rightarrow M$ and $A^* \rightarrow M$ be any dual vector bundles and let $\langle \langle \cdot, \cdot \rangle \rangle$ denote the pairing of $T^2(A^*)$ and $T^2(A)$ over T^2M used in the proof above. Then, for $\Psi \in T^2(A^*)$ and $\xi \in T^2(A)$ projecting to the same element of T^2M ,*

$$\langle \langle J_{A^*}(\Psi), J_A(\xi) \rangle \rangle = \langle \langle \Psi, \xi \rangle \rangle.$$

PROOF. Write $\xi = \partial_s \partial_t a(0,0)$ for a smooth $a(s,t) \in A$. The notation means that a is defined in a neighbourhood of the origin, and that ξ is the tangent vector at 0 to the curve $s \mapsto \partial_t a(s,0) = \frac{\partial a}{\partial t}(s,0)$ in TM .

The projection condition ensures that we can likewise write $\Psi = \partial_s \partial_t b(0,0)$ for a smooth patch $b(s,t) \in A^*$ with $q(a(s,t)) = q_*(b(s,t))$ for all (s,t) near $(0,0)$. Now

$$\langle \langle \Psi, \xi \rangle \rangle = \left. \frac{d}{ds} \langle \partial_t a(s,0), \partial_t b(s,0) \rangle \right|_{s=0} = \left. \frac{\partial^2}{\partial s \partial t} \langle a(s,t), b(s,t) \rangle \right|_{(0,0)}$$

and the result follows from the equality of mixed partials. $\square$

We now return to the proof of 4.3. From Lemma 4.5 and (22) we have

$$\pi_{TA}^\# \circ R_\bullet = J_A \circ T(\pi_A^\# \circ R_A) \circ \Theta_{A^*}^{-1} \circ I_A^\times.$$

Since (A, A^*) is a Lie bialgebroid, $\pi_A^\# \circ R_A: T^*A^* \rightarrow TA$ is a morphism of Lie algebroids over a_* , so $T(\pi_A^\# \circ R_A)$ is a morphism of the prolongation structures over $T(a_*)$. Also, since I_A is a Poisson diffeomorphism, $I_A^\times$ is an isomorphism of the cotangent Lie algebroids over I_A^{-1} .

For any Poisson manifold, $\Theta_P: T(T^*P) \rightarrow T^*(TP)$ is an isomorphism of Lie algebroids over TP from the tangent prolongation of the cotangent Lie algebroid structure on T^*P to the cotangent Lie algebroid of the tangent Poisson structure [15, 10.3.13]. We apply this to $P = A^*$.

Finally we apply Lemma 4.7(i). Altogether, we have that $\pi_{TA}^\# \circ R_\bullet$ is a Lie algebroid morphism and this completes the proof of Proposition 4.3. $\square$

Lemma 4.7 (i) *Let A be a Lie algebroid on M and let $T^2A \rightarrow T^2M$ be the iterated prolongation Lie algebroid. Then the canonical involution $J_A: T^2A \rightarrow T^2A$ is a Lie algebroid automorphism over J_M .*

(ii) *For any Poisson manifold P , the canonical involution J_P is a Poisson diffeomorphism, with respect to the iterated tangent lift Poisson structure.*

PROOF. (i) It is sufficient to show that the dual map $J_A^* T^2M$ is a Poisson diffeomorphism. The dual of $T^2A \rightarrow T^2M$ is $T^\bullet(TA) \rightarrow T^2M$ and this is isomorphic to $T(T^\bullet A) \rightarrow T^2M$ under the map I for $TA \rightarrow TM$. Composing with $T(I_A)$ we have that $T^\bullet(TA) \rightarrow T^2M$ is isomorphic to $T^2(A^*) \rightarrow T^2M$ and with this identification, J_A becomes $J_{A^*}: T^2(A^*) \rightarrow T^2(A^*)$. The result now follows from (ii).

(ii) Write $J = J_P$ and $J_T = J_{TP}$. We must show that $T(J) \circ \pi_{T^2P}^\# = \pi_{T^2P}^\# \circ J^\times$. Using the definition of $\pi_{T^2P}^\#$, we have to show that

$$T(J) \circ J_T \circ T(J) \circ J_T \circ T(J) \circ T^2(\pi^\#) \circ T(\Theta)^{-1} \circ \Theta_T^{-1} = T^2(\pi^\#) \circ T(\Theta)^{-1} \circ \Theta_T^{-1} \circ J^\times.$$

Now $T(J) \circ J_T \circ T(J) \circ J_T \circ T(J) = J_T$. If the coordinates in $\mathbb{R}^3$ are t_1, t_2, t_3 then $T(J)$ corresponds to the permutation (2 3) and J_T to the permutation (1 2). Using this and the naturality property $J_T \circ T^2(\pi^\#) = T^2(\pi^\#) \circ J_{T^*P}$, we are reduced to showing that, for any manifold P ,

$$J_P^\times \circ \Theta_{TP} \circ T(\Theta_P) = \Theta_{TP} \circ T(\Theta_P) \circ J_{T^*P}. \quad (26)$$

Take $\xi \in T^2(T^*P)$ and $\zeta \in T^3P$ with $T^2(c)(\xi) = p_{Y^2}(\zeta)$. Then, using 4.4,

$$\langle \Theta_T \circ T(\Theta) \circ J_{T^*P}(\xi), \zeta \rangle = \langle T(\Theta) \circ J_{T^*P}(\xi), J_T(\zeta) \rangle$$

and

$$\langle J^\times \circ \Theta_T \circ T(\Theta)(\xi), \zeta \rangle = \langle T(\Theta)(\xi), J_T \circ T(J)(\zeta) \rangle$$

and equality follows as in Lemma 4.6. $\square$

Returning to the proof of 4.1, suppose that A is a Lie algebroid on M and that A^* has a Lie algebroid structure, not *a priori* related to the structure on A . Consider $D = T^*A$ with the two cotangent Lie algebroid structures arising from the Poisson structures on A^* and A , and suppose that these structures make D a double Lie algebroid.

Then in particular, by Theorem 3.9 the anchor $T^*A \rightarrow TA^*$ of the horizontal structure must be a Lie algebroid morphism over $a: A \rightarrow TM$ with respect to the vertical structures. The anchor is

$$T^*A \xrightarrow{R_{A^*}} T^*A^* \xrightarrow{\pi_{A^*}^\#} TA^*$$

That this be a Lie algebroid morphism over a is precisely the dual form of 3.3.

This completes the proof of Theorem 4.1. $\square$

Recall the Manin triple characterization of a Lie bialgebra, as given in [14]: Given a Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$ the vector space direct sum $\mathfrak{d} = \mathfrak{g} \oplus \mathfrak{g}^*$ has a Lie algebra bracket defined in terms of the two coadjoint representations; this is the classical Drinfel'd double of $(\mathfrak{g}, \mathfrak{g}^*)$. This bracket is invariant under the pairing $\langle X + \varphi, Y + \psi \rangle = \langle \psi, X \rangle + \langle \varphi, Y \rangle$ and both $\mathfrak{g}$ and $\mathfrak{g}^*$ are coisotropic subalgebras. Conversely, if a Lie algebra $\mathfrak{d}$ is a vector space direct sum $\mathfrak{g} \oplus \mathfrak{h}$, both of which are coisotropic with respect to an invariant pairing of $\mathfrak{d}$ with itself, then $\mathfrak{h} \cong \mathfrak{g}^*$ and $(\mathfrak{g}, \mathfrak{h})$ is a Lie bialgebra, with $\mathfrak{d}$ as the double.

This result provided a characterization of the notion of Lie bialgebra in terms of a single Lie algebra structure on $\mathfrak{d}$, the conditions being expressed in terms of the simple notion of pairing. It also demonstrated, as a simple consequence, that the notion of Lie bialgebra is self-dual: $(\mathfrak{g}, \mathfrak{g}^*)$ is a Lie bialgebra if and only if $(\mathfrak{g}^*, \mathfrak{g})$ is so.

For Lie bialgebroids, Kosmann–Schwarzbach [9] gave an elegant proof of self-duality, in terms of her criterion that (A, A^*) is a Lie bialgebroid if and only if the coboundary of one structure is a derivation of the Schouten bracket of the other.

For a Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$, the structure of the cotangent double Lie algebroid $T^*\mathfrak{g} \cong \mathfrak{g} \times \mathfrak{g}^*$ equips $T^*\mathfrak{g}$ with Lie algebroid structures on bases $\mathfrak{g}$ and $\mathfrak{g}^*$. These are the action Lie algebroids defined by the coadjoint actions of $\mathfrak{g}$ and $\mathfrak{g}^*$ on each other. We show in §5 that a double Lie algebroid $(D; A, B.M)$ for which the core is trivial — as is the case for $D = T^*\mathfrak{g}$ — has a third Lie algebroid structure, on base M . In the case of $T^*\mathfrak{g}$, M is a point and this third structure is the Drinfel’d double Lie algebra.

Despite the differences, it seems legitimate to regard Theorem 4.1 as a generalization of the Manin triple theorem, since it provides a criterion, in terms of a notion of double, for Lie algebroid structures on a pair of dual bundles to constitute a Lie bialgebroid.

In [11], Liu, Weinstein and Xu place a bracket structure on $A \oplus A^*$, where (A, A^*) is a Lie bialgebroid, which is a direct generalization of the Lie algebra structure of the Drinfel’d double, and they prove a direct generalization of the Manin triple theorem. This is not a Lie algebroid structure, but a Courant algebroid, a vector bundle with a bracket structure in which the Jacobi and Leibniz identities do not necessarily hold. Whereas the cotangent double has two simple structures which interact in a nontrivial way, the single bracket structure on $A \oplus A^*$ of [11] is less well-behaved but embodies the whole structure.

The notion of Courant algebroid has been developed in work on Dirac structures and generalized complex geometry. On the other hand, the notion of double Lie algebroid abstracts iterated and second-order constructions in differential geometry (see particularly §6), and provides a general setting for duality phenomena. In particular, there is a clear global form of the notion of double Lie algebroid.

Another important property of the classical Drinfel’d double is that $\mathfrak{g} \bowtie \mathfrak{g}^*$ itself has a natural structure of Lie bialgebra. A double form of this result will be given in [25].

§ 5 MATCHED PAIRS AND VACANT DOUBLE LIE ALGEBROIDS

A brief history of matched pairs of Lie algebras was given in the Introduction. The corresponding concept of a matched pair of Lie groups [14], [28] was extended to Lie groupoids in [16]. In [31], Mokri differentiated the twisted automorphism equations of [16] to obtain conditions on a pair of Lie algebroid representations, of A on B and of B on A , which ensure that the direct sum vector bundle $A \oplus B$ has a Lie algebroid structure with A and B as subalgebroids. We quote the following.

Definition 5.1 [31, 4.2] *Let A and B be Lie algebroids on base M , with anchors a and b , and let $\rho: A \rightarrow \mathfrak{D}(B)$ and $\sigma: B \rightarrow \mathfrak{D}(A)$ be representations of A on the vector bundle B and of B on the vector bundle A . Then A and B together with ρ and σ form a matched pair of Lie algebroids if equations (1) from the Introduction hold.*

The Lie algebroid $\mathfrak{D}(E)$ of derivations on E is defined in §2. The proof of the following proceeds along the same lines as the proof for Lie algebras.

Proposition 5.2 [31, 4.3] *Given a matched pair of Lie algebroids, there is a Lie algebroid structure $A \bowtie B$ on the direct sum vector bundle $A \oplus B$, with anchor $c(X \oplus Y) = a(X) + b(Y)$ and bracket*

$$[X_1 \oplus Y_1, X_2 \oplus Y_2] = \{[X_1, X_2] + \sigma_{Y_1}(X_2) - \sigma_{Y_2}(X_1)\} \oplus \{[Y_1, Y_2] + \rho_{X_1}(Y_2) - \rho_{X_2}(Y_1)\}.$$

Conversely, if $A \oplus B$ has a Lie algebroid structure for which $A \oplus 0$ and $0 \oplus B$ are Lie subalgebroids, then ρ and σ defined by $[X \oplus 0, 0 \oplus Y] = -\sigma_Y(X) \oplus \rho_X(Y)$ form a matched pair.

We now show that matched pairs of Lie algebroids correspond precisely to double Lie algebroids for which the core is zero.

Definition 5.3 *A double Lie algebroid $(D; A, B; M)$ is vacant if the core is the zero bundle.*

Equivalently, a double Lie algebroid is vacant if the combination of the two projections, $(q_A^D, q_B^D): D \rightarrow A \times_M B$ is a diffeomorphism. This is a direct analogue of the definition of vacancy for double Lie groupoids and for $\mathcal{L}\mathcal{A}$ -groupoids [16, 2.11, 4.10].

Consider a vacant double Lie algebroid $(D; A, B; M)$. Note that $D \rightarrow A$ and $D \rightarrow B$ are the pullback bundles $q_A^!B$ and $q_B^!A$. The dual $D^*A \rightarrow C^*$ is the Whitney sum $A \oplus B^*$. The duality between D^*A and D^*B is

$$\langle X + \psi, \varphi + Y \rangle = \langle \psi, Y \rangle - \langle \varphi, X \rangle. \quad (27)$$

The horizontal bundle projection $q_B^D: D \rightarrow B$ is a morphism of Lie algebroids over $q_A: A \rightarrow M$ and since it is a fibrewise surjection, it defines an action of B on q_A as in [15, §4.1]. Namely, each section Y of B induces the pullback section $1 \otimes Y$ of $q_A^!B$ and this induces a vector field $\eta(Y) = \Delta_A(1 \otimes Y)$ on A , where $\Delta_A: D \rightarrow TA$ is the anchor of the vertical structure. Since the anchor is a morphism of double vector bundles, $\eta(Y)$ is linear over the vector field $b(Y)$ on M , in the sense of [15, §3.4]; that is, $\eta(Y)$ is a vector bundle morphism $A \rightarrow TA$ over $b(Y): M \rightarrow TM$. It follows that $\eta(Y)$ defines derivations $\sigma_Y^{(*)}$ on A^* and σ_Y on A by

$$\eta(Y)(\ell_\varphi) = \ell_{\sigma_Y^{(*)}(\varphi)}, \quad \langle \varphi, \sigma_Y(X) \rangle = b(Y)\langle \varphi, X \rangle - \langle \sigma_Y^{(*)}(\varphi), X \rangle \quad (28)$$

where $\varphi \in \Gamma A^*$, $X \in \Gamma A$, and ℓ_φ denotes the function $A \rightarrow \mathbb{R}$, $X \mapsto \langle \varphi(q_A X), X \rangle$. Since q_A^D is a Lie algebroid morphism, it follows that σ is a representation of B on the vector bundle A , with contragredient representation $\sigma^{(*)}$. We will use both σ and $\sigma^{(*)}$ in what follows.

Likewise, q_A^D is a morphism of Lie algebroids over q_B and for each $X \in \Gamma A$ we obtain a linear vector field $\xi(X) \in \mathcal{X}(B)$ over $a(X)$. We define derivations $\rho_X^{(*)}$ on B^* and ρ_X on B by

$$\xi(X)(\ell_\psi) = \ell_{\rho_X^{(*)}(\psi)}, \quad \langle \psi, \rho_X(Y) \rangle = a(X)\langle \psi, Y \rangle - \langle \rho_X^{(*)}(\psi), Y \rangle. \quad (29)$$

Again, $\rho^{(*)}$ and ρ are representations of A .

The two Lie algebroid structures on D are action Lie algebroids determined by the actions $Y \mapsto \eta(Y)$ and $X \mapsto \xi(X)$ [15, §4.1]. It therefore follows that the dual Poisson structures are semi-direct, but we need to prove this directly. We first calculate the Lie algebroid structure on $A^* \oplus B$ induced from that on $D \rightarrow A$.

Lemma 5.4 *The Lie algebroid structure on $D \star A \star C^* = A^* \oplus B$ induced from the double Lie algebroid structure has anchor $e(\varphi \oplus Y) = b(Y)$ and bracket*

$$[\varphi_1 \oplus Y_1, \varphi_2 \oplus Y_2] = \{\sigma_{Y_1}^{(*)}(\varphi_2) - \sigma_{Y_2}^{(*)}(\varphi_1)\} \oplus [Y_1, Y_2]. \quad (30)$$

The Lie algebroid structure on $D \star A = A \oplus B^$ induced from the double Lie algebroid structure has anchor $e_*(X \oplus \psi) = -a(X)$ and bracket*

$$[X_1 \oplus \psi_1, X_2 \oplus \psi_2] = [X_2, X_1] \oplus \{\rho_{X_2}^{(*)}(\psi_1) - \rho_{X_1}^{(*)}(\psi_2)\}. \quad (31)$$

PROOF. We apply equations (19). Firstly, a linear section (ξ, Y) of $D \rightarrow A$ is necessarily a pullback section $\xi(X) = X \oplus Y(m)$ for $X \in A_m$. From the definition (13) of $\xi^\square$ we find that $\xi^\square \in \Gamma(A^* \oplus B)$ is

$$\xi^\square(m) = 0_m^{A^*} \oplus Y(m).$$

It follows from the first equation in (19) that $[0 \oplus Y_1, 0 \oplus Y_2] = 0 \oplus [Y_1, Y_2]$ for $Y_1, Y_2 \in \Gamma B$.

Next, $\varphi \in \Gamma A^*$ defines $\overline{\varphi} \in \Gamma(A^* \oplus B)$ where $\overline{\varphi}(m) = \varphi(m) \oplus 0_m^B$. It follows from (19) that $[\varphi_1 \oplus 0, \varphi_2 \oplus 0] = 0 \oplus 0$.

The second equation in (19) is $[\xi^\square, \overline{\varphi}] = \overline{\Lambda_\xi(\varphi)}$. Comparing (17) with (28), we have $\Lambda = \sigma^{(*)}$. So $[0 \oplus Y, \varphi \oplus 0] = \sigma_Y^{(*)}(\varphi) \oplus 0$ and this completes the proof of (30).

For (31) apply (19) to $D \star B \star C^* = A \oplus B^*$ and transport the structure to $D \star A = A \oplus B^*$ by $Z_A(X, \psi) = (-X, \psi)$. $\square$

Thus $A^* \oplus B$ is the semi-direct product $A^* \rtimes B$ of B with the vector bundle A^* with respect to $\sigma^{(*)}$. (This is the simple semi-direct product of two Lie algebroids on the same base, as in [15, p.270].) We denote the structure on $A \oplus B^*$ by $A^{op} \rtimes B^*$.

Theorem 5.5 *Let $(D; A, B; M)$ be a vacant double Lie algebroid. Then the two Lie algebroid structures on D are action Lie algebroids corresponding to actions which define representations ρ , of A on B , and σ , of B on A , with respect to which A and B form a matched pair.*

Conversely, let A and B be a matched pair of Lie algebroids with respect to representations ρ and σ . Then the action of A on q_B induced by ρ and the action of B on q_A induced by σ define Lie algebroid structures on $D = A \times_M B$ with respect to which $(D; A, B; M)$ is a vacant double Lie algebroid.

PROOF. We first need to prove the three equations (1). We start by applying the bialgebroid condition to $E = A^* \rtimes B$ and $E^* = A^{op} \rtimes B^*$. We use the dual form of (20), which in this case is :

$$d^{E^*}[\varphi_1 \oplus Y_1, \varphi_2 \oplus Y_2] = [d^{E^*}(\varphi_1 \oplus Y_1), \varphi_2 \oplus Y_2] + [\varphi_1 \oplus Y_1, d^{E^*}(\varphi_2 \oplus Y_2)] \quad (32)$$

for all $\varphi_1 \oplus Y_1, \varphi_2 \oplus Y_2 \in \Gamma E$. By replacing $\varphi_2 \oplus Y_2$ by $f\varphi_2 \oplus fY_2$, for $f \in C^\infty(M)$, and expanding out, we find, as in [15, p.449], that

$$d^{E^*}[\varphi \oplus Y, f] = [d^{E^*}(\varphi \oplus Y), f] + [\varphi \oplus Y, d^{E^*}(f)]. \quad (33)$$

Now $d^{E^*}(f) = -d^A f \oplus 0$. So, for the last term, we have $[\varphi \oplus Y, d^{E^*}(f)] = -\sigma_Y^{(*)}(d^A f) \oplus 0$, and pairing this with any $X \oplus \psi \in \Gamma E^*$. we have

$$\langle [\varphi \oplus Y, d^{E^*}(f)], X \oplus \psi \rangle = a(\sigma_Y(X))(f) - b(Y)(a(X)(f)).$$

On the LHS, $[\varphi \oplus Y, f] = e(\varphi \oplus Y)(f) = b(Y)(f)$ and so, pairing with $X \oplus \psi$ gives

$$\langle d^{E*}(b(Y)(f)), X \oplus \psi \rangle = e_*(X \oplus \psi)(b(Y)(f)) = -a(X)(b(Y)(f)).$$

Consider the first term on the RHS of (33). For any $\xi \in \Gamma\Lambda^2(E)$ we have $[\xi, f] = -\iota_{df}(\xi)$, [15, 12.1.4], so pairing with $X \oplus \psi$ gives

$$\langle [\xi, f], X \oplus \psi \rangle = -\xi(df, X \oplus \psi).$$

We need the following lemma, which is a straightforward calculation.

Lemma 5.6

$$d^{E*}(\varphi \oplus Y)(X_1 \oplus \psi_1, X_2 \oplus \psi_2) = -(d^A\varphi)(X_1, X_2) + \langle \psi_1, \rho_{X_2}(Y) \rangle - \langle \psi_2, \rho_{X_1}(Y) \rangle.$$

□

For the first term on the RHS of (33), we therefore have

$$\begin{aligned} \langle [d^{E*}(\varphi \oplus Y), f], X \oplus \psi \rangle &= -d^{E*}(\varphi \oplus Y)(d^E f, X \oplus \psi) = \\ &= -\langle d^B f, \rho_X(Y) \rangle = -b(\rho_X(Y))(f). \end{aligned}$$

using $d^E f = 0 \oplus d^B f$. Putting these together, (33) becomes

$$-a(X)(b(Y)(f)) = -b(\rho_X(Y))(f) + a(\sigma_Y(X))(f) - b(Y)(a(X)(f)),$$

which proves the third equation in (1).

Now return to the bialgebroid equation (32). Set $\varphi_1 = \varphi_2 = 0$, so that we have

$$d^{E*}[0 \oplus Y_1, 0 \oplus Y_2] = [d^{E*}(0 \oplus Y_1), 0 \oplus Y_2] + [0 \oplus Y_1, d^{E*}(0 \oplus Y_2)] \quad (34)$$

and evaluate this at $(0 \oplus \psi) \wedge (X \oplus 0)$. The LHS of (34) is easily seen to be $\langle \psi, \rho_X[Y_2, Y_1] \rangle$.

On the RHS, consider the second term first. Writing the bracket as the Lie derivative of the multisection $d^{E*}(0 \oplus Y_2)$ with respect to $0 \oplus Y_1$, we have

$$\begin{aligned} &\langle \mathfrak{L}_{0 \oplus Y_1}(d^{E*}(0 \oplus Y_2)), (0 \oplus \psi) \wedge (X \oplus 0) \rangle \\ &= \mathfrak{L}_{0 \oplus Y_1} \langle d^{E*}(0 \oplus Y_2), (0 \oplus \psi) \wedge (X \oplus 0) \rangle - \langle d^{E*}(0 \oplus Y_2), \mathfrak{L}_{0 \oplus Y_1}((0 \oplus \psi) \wedge (X \oplus 0)) \rangle \end{aligned} \quad (35)$$

In the first term, $d^{E*}(0 \oplus Y_2)(0 \oplus \psi, X \oplus 0) = \langle \psi, \rho_X(Y_2) \rangle$, and $e(0 \oplus Y_1) = b(Y_1)$, so we have $b(Y_1)\langle \psi, \rho_X(Y_2) \rangle$.

For the second term, note first that $\mathfrak{L}_{0 \oplus Y}(0 \oplus \psi) = 0 \oplus \mathfrak{L}_Y(\psi)$ and $\mathfrak{L}_{0 \oplus Y}(X \oplus 0) = \sigma_Y(X) \oplus 0$. Using these, the second term on the RHS of (35) is

$$\langle \mathfrak{L}_{Y_1}(\psi), \rho_X(Y_2) \rangle + \langle \psi, \rho_{\sigma_{Y_1}(X)}(Y_2) \rangle.$$

Calculating the first term on the RHS of (34) in the same way, we have

$$\begin{aligned} \langle \psi, \rho_X[Y_2, Y_1] \rangle &= -b(Y_2)\langle \psi, \rho_X(Y_1) \rangle + \langle \mathfrak{L}_{Y_2}(\psi), \rho_X(Y_1) \rangle + \langle \psi, \rho_{\sigma_{Y_2}(X)}(Y_1) \rangle \\ &\quad + b(Y_1)\langle \psi, \rho_X(Y_2) \rangle - \langle \mathfrak{L}_{Y_1}(\psi), \rho_X(Y_2) \rangle - \langle \psi, \rho_{\sigma_{Y_1}(X)}(Y_2) \rangle. \end{aligned}$$

Finally, $b(Y_1)\langle\psi, \rho_X(Y_2)\rangle - \langle\mathfrak{L}_{Y_1}(\psi), \rho_X(Y_2)\rangle = \langle\psi, [Y_1, \rho_X(Y_2)]\rangle$ and we have the first equation of (1). The second equation is proved in the same way.

Now consider the second part of 5.5. It is straightforward to prove that $(A \times_M B; A, B; M)$, with the action Lie algebroid structures on $A \times_M B$, is an $\mathcal{L}\mathcal{A}$ -vector bundle both horizontally and vertically. We must verify the bialgebroid condition (32).

Firstly, (34) at $(0 \oplus \psi) \wedge (X \oplus 0)$ can be proved by reversing the argument above. The case of any $(X_1 \oplus \psi_1, X_2 \oplus \psi_2) = (X_1 \oplus 0, 0 \oplus \psi_2) + (0 \oplus \psi_1, X_2 \oplus 0)$ follows from the skew-symmetry of all the terms. Thus (34) holds.

For (32) with $Y_1 = Y_2 = 0$ and any arguments, it is straightforward to see that all terms are identically zero. It remains to consider (32) with $\varphi_1 = 0, Y_2 = 0$. For the LHS we have

$$d^{E^*}[0 \oplus Y, \varphi \oplus 0] = -d^A(\sigma_Y^{(*)}(\varphi)) \oplus 0.$$

Expanding out $-d^A(\sigma_Y^{(*)}(\varphi))(X_1, X_2)$ we have

$$\begin{aligned} & -a(X_1)b(Y)\langle\varphi, X_2\rangle + a(X_1)\langle\varphi, \sigma_Y(X_2)\rangle + a(X_2)b(Y)\langle\varphi, X_1\rangle \\ & -a(X_2)\langle\varphi, \sigma_Y(X_1)\rangle + b(Y)\langle\varphi, [X_1, X_2]\rangle - \langle\varphi, \sigma_Y([X_1, X_2])\rangle. \end{aligned} \quad (36)$$

On the RHS the second term is $[0 \oplus Y, d^{E^*}(\varphi \oplus 0)] = [0 \oplus Y, -d^A\varphi \oplus 0] = -\sigma_Y^{(*)}(d^A\varphi) \oplus 0$, using 5.6 and (30), and extending $\sigma_Y^{(*)}$ to the exterior square. Expanding out $\langle-\sigma_Y^{(*)}(d^A\varphi), X_1 \wedge X_2\rangle$, we have

$$\begin{aligned} & -b(Y)a(X_1)\langle\varphi, X_2\rangle + b(Y)a(X_2)\langle\varphi, X_1\rangle + b(Y)\langle\varphi, [X_1, X_2]\rangle \\ & + a(\sigma_Y(X_1))\langle\varphi, X_2\rangle - a(X_2)\langle\varphi, \sigma_Y(X_1)\rangle - \langle\varphi, [\sigma_Y(X_1), X_2]\rangle \\ & + a(X_1)\langle\varphi, \sigma_Y(X_2)\rangle - a(\sigma_Y(X_2))\langle\varphi, X_1\rangle - \langle\varphi, [X_1, \sigma_Y(X_2)]\rangle. \end{aligned} \quad (37)$$

The first term on the RHS, evaluated at $F_1 \wedge F_2 \in \Gamma\Lambda^2(E^*)$, is

$$\begin{aligned} \langle[d^{E^*}(0 \oplus Y), \varphi \oplus 0], F_1 \wedge F_2\rangle &= \langle-\mathfrak{L}_{\varphi \oplus 0}(d^{E^*}(0 \oplus Y)), F_1 \wedge F_2\rangle \\ &= \langle d^{E^*}(0 \oplus Y), \mathfrak{L}_{\varphi \oplus 0}(F_1 \wedge F_2)\rangle, \end{aligned} \quad (38)$$

since $e(\varphi \oplus 0) = 0$. Now $\mathfrak{L}_{\varphi \oplus 0}(X \oplus \psi) = 0 \oplus \theta_{\varphi, X}$ where $\theta_{\varphi, X} \in B^*$ is the element such that

$$\langle Z, \theta_{\varphi, X}\rangle = \langle\sigma_Z^{(*)}(\varphi), X\rangle$$

for all $Z \in B$. Put $F_i = X_i \oplus \psi_i$ in (38) and write $\theta_i = \theta_{\varphi, X_i}$. Then (38) becomes

$$\begin{aligned} & \langle d^{E^*}(0 \oplus Y), (0 \oplus \theta_1) \wedge (X_2 \oplus \psi_2)\rangle + \langle d^{E^*}(0 \oplus Y), (X_1 \oplus \psi_1) \wedge (0 \oplus \theta_2)\rangle \\ &= a(X_2)\langle Y, \theta_1\rangle - \langle 0 \oplus Y, 0 \oplus \rho_{X_2}^{(*)}(\theta_1)\rangle - a(X_1)\langle Y, \theta_2\rangle + \langle 0 \oplus Y, 0 \oplus \rho_{X_1}^{(*)}(\theta_2)\rangle \\ &= b(\rho_{X_2}(Y))\langle\varphi, X_1\rangle - \langle\varphi, \sigma_{\rho_{X_2}(Y)}(X_1)\rangle - b(\rho_{X_1}(Y))\langle\varphi, X_2\rangle + \langle\varphi, \sigma_{\rho_{X_1}(Y)}(X_2)\rangle. \end{aligned}$$

Adding this to (37) gives (36), using the first and last equations of (1). The proof of Theorem 5.5 is complete. $\square$

In the process we have also proved the following result.

Corollary 5.7 *Let A and B be Lie algebroids on the same base M , and let $\rho: A \rightarrow \mathfrak{D}(B)$ and $\sigma: B \rightarrow \mathfrak{D}(A)$ be representations on the vector bundles underlying B and A .*

Then A , B , ρ , σ form a matched pair if and only if $(A^ \rtimes B, A^{op} \rtimes B^*)$, with the Lie algebroid structures described in 5.4, is a Lie bialgebroid.*

Thus the three matched pair equations in (1) are embodied in the bialgebroid equation (32). Theorem 5.5 actually shows the equivalence of three formulations: a vacant double Lie algebroid structure with sides A and B is equivalent to a matched pair structure on (A, B) , which in turn is equivalent to a Lie bialgebroid structure on $(A^* \rtimes B, A^{op} \rtimes B^*)$ of the type in 5.4. We thus have a diagrammatic characterization of matched pairs of Lie algebroids: recall that the characterization 3.3 of a Lie bialgebroid is formulated entirely in terms of the Poisson tensor and the canonical isomorphism R . Thus 5.5 provides a definition of matched pair which can be formulated more generally in a category possessing pullbacks and suitable additive structure. The corresponding characterization of matched pairs of Lie groupoids in terms of vacant double Lie groupoids is in [16, §2].

Given a vacant double Lie algebroid $(D; A, B; M)$ the Lie algebroid structure $A \bowtie B$ on $D \rightarrow M$ may be called a *diagonal structure* by analogy with the groupoid case.

Forms of Corollary 5.7 in the Lie algebra case have been found by Majid [29, §8.3] and in the setting of Lie–Rinehart algebras by Huebschmann [7].

In the case of a Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$, the Lie bialgebroid associated to the vacant double Lie algebroid is $(\mathfrak{g} \oplus \mathfrak{g}_0, \mathfrak{g}^* \oplus \mathfrak{g}_0^*)$ where the subscripts denote the abelianizations. This is of course consistent with 4.1 in the bialgebra case — which is both a bialgebroid and a matched pair.

§ 6 THE DOUBLE LIE ALGEBROID OF A DOUBLE LIE GROUPOID

Definition 3.1 of a double Lie algebroid arose from work on the double Lie algebroid of a double Lie groupoid, as constructed in [16], [20]. Some of what is required in order to verify that the double Lie algebroid of a double Lie groupoid does satisfy 3.1 has been given in [18], and we recall the details briefly.

First we recall the notion of double Lie groupoid (see [16] and references given there). A double Lie groupoid consists of a manifold S equipped with two Lie groupoid structures on bases H and V , each of which is a Lie groupoid on base M , such that the structure maps (source, target, multiplication, identity, inversion) of each groupoid structure on S are morphisms with respect to the other, and such that the *double source map* $\alpha_2: S \rightarrow H \times_M V$, $s \mapsto (\tilde{\alpha}_V(s), \tilde{\alpha}_H(s))$ is a surjective submersion; see Figure 7(a). One should think of elements of S as squares, the horizontal edges of which come from H , the vertical edges from V , and the corner points from M .

Consider a double Lie groupoid $(S; H, V; M)$ as in Figure 7(a). Applying the Lie functor to the vertical structure $S \rightrightarrows H$ gives a Lie algebroid $A_V S \rightarrow H$ which also has a groupoid structure over AV obtained by applying the Lie functor to the structure maps of $S \rightrightarrows V$; this is the *vertical $\mathcal{LA}$ -groupoid* of S [16, §4], as in Figure 7(b). We recall the notion of $\mathcal{LA}$ -groupoid from [16].

Definition 6.1 *An $\mathcal{LA}$ -groupoid is a manifold Ω equipped with a Lie groupoid structure on a base A , which is a Lie algebroid on a base M , and a Lie algebroid structure on a base G ,*




Figure 7.

which is a Lie groupoid on M , such that the groupoid structure maps (source, target, identity, division) are morphisms of Lie algebroids, and the map $\Omega \rightarrow G \times_M A$ which combines the bundle projection and the Lie groupoid source, is a surjective submersion.

The Lie algebroid of $A_V S \rightrightarrows AV$ is denoted $A^2 S$; there is a double vector bundle structure $(A^2 S; AH, AV; M)$ obtained by applying A to the vector bundle structure of $A_V S \rightarrow H$ [20]; see Figure 7(c). Reversing the order of these operations, one defines first the *horizontal* $\mathcal{LA}$ -groupoid $(A_H S; AH, V; M)$ and then takes the Lie algebroid $A_2 S = A(A_H S)$. The canonical involution $J_S: T^2 S \rightarrow T^2 S$ then restricts to an isomorphism of double vector bundles $\tilde{J}_S: A^2 S \rightarrow A_2 S$ and allows the Lie algebroid structure on $A^2 S \rightarrow AV$ to be transported to $A_2 S \rightarrow AV$. Thus $A_2 S$ is a double vector bundle equipped with four Lie algebroid structures; in [20] we called this the *double Lie algebroid of S* . The core of both double vector bundles $A_2 S$ and $A^2 S$ is AK , the Lie algebroid of the core groupoid $K \rightrightarrows M$ of S [20, 1.6].

Theorem 6.2 *With the structures just defined, $(A_2 S; AH, AV; M)$ is a double Lie algebroid.*

PROOF. Write $D = A_2 S$. The structure maps for the horizontal vector bundle $A_2 S \rightarrow AV$ are obtained by applying the Lie functor to the structure maps of $A_H S \rightarrow V$ and are therefore Lie algebroid morphisms with respect to the vertical Lie algebroid structures. The corresponding statement is true for the vertical vector bundle $A^2 S \rightarrow AH$ and this is transported by $\tilde{J}_S$ to $A_2 S \rightarrow AH$. Thus the first condition of 3.1 is satisfied.

Now consider the bialgebroid condition. We first need to identify the Lie algebroid structures on $A_2 S \star AH \star A^* K$ and $A_2 S \star AV \star A^* K$. From [18] we know that $A_V^* S \rightrightarrows A^* K$ is a Poisson groupoid, and therefore $(A(A_V^* S), A^*(A_V^* S))$ is a Lie bialgebroid. To relate these structures, we need to recall the duality of $\mathcal{VB}$ -groupoids.

Consider any $\mathcal{VB}$ -groupoid as in Figure 8(a), with core vector bundle $L \rightarrow M$. Taking the dual in the sense of $\mathcal{VB}$ -groupoids [32], [15, §11.2], we have the $\mathcal{VB}$ -groupoid $(\Omega^*; G, L^*; M)$ as in Figure 8(b).

The pairing of Ω and Ω^* is a Lie groupoid morphism $\Omega \times_G \Omega^* \rightarrow \mathbb{R}$ and the induced pairing $\langle\langle \cdot, \cdot \rangle\rangle = A(\langle \cdot, \cdot \rangle)$ of the Lie algebroids is also non-degenerate and respects the double vector bundle structures [15, §11.5]. It therefore induces an isomorphism of double vector bundles

$$\mathfrak{I}_\Omega: A(\Omega^*) \rightarrow A\Omega \star AG$$

which preserves both side bundles and the core. Write $\tilde{I}_V$ and $\tilde{I}_H$ for this map when $\Omega = A_V S$ and $\Omega = A_H S$. Composing the inverse of $\tilde{J}_S \star AH$ with $\tilde{I}_V$ gives an isomorphism

$$\tilde{\Theta}_H = (\tilde{J}_S \star AH)^{-1} \circ \tilde{I}_V: A(A_V^* S) \rightarrow A_2 S \star AH. \quad (39)$$




Figure 8.

Dualizing this over A^*K we get $\tilde{\Theta}_H \star A^*K: A_2S \star AH \star A^*K \rightarrow A^*(A_V^*S)$ and to prove that this is an isomorphism of Lie algebroids it is sufficient to prove that $\tilde{\Theta}_H$ is a Poisson map.

On the other hand, consider $\tilde{I}_H: A(A_H^*S) \rightarrow A_2S \star AV$. Dualizing this over A^*K we obtain

$$\tilde{I}_H \star A^*K: A_2S \star AV \star A^*K \rightarrow A^*(A_H^*S).$$

Again, to show that this is an isomorphism of Lie algebroids, it is sufficient to show that $\tilde{I}_H$ is a Poisson map. However the Poisson structure on $A_2S \star AV$ is the dual of the Lie algebroid structure on $A_2S \rightarrow AH$ and this is obtained from the Lie algebroid structure on $A^2S \rightarrow AH$, transported by $\tilde{J}_S$. So we actually need to show that $\tilde{\Theta}_V$, defined by

$$\tilde{\Theta}_V = (\tilde{J}_S \star AV) \circ \tilde{I}_H: A(A_H^*S) \rightarrow A^2S \star AV \quad (40)$$

is a Poisson map. The proof is therefore reduced to the following.

Proposition 6.3 *The map $\tilde{\Theta}_H: A(A_V^*S) \rightarrow A_2S \star AH$ is a Poisson map with respect to the Poisson structure on $A(A_V^*S)$ induced from the Poisson structure on A_V^*S which is dual to the Lie algebroid $A_VS \rightarrow H$, and the Poisson structure on $A_2S \star AH$ dual to the Lie algebroid $A_2S \rightarrow AH$.*

*The map $\tilde{\Theta}_V: A(A_H^*S) \rightarrow A^2S \star AV$ is a Poisson map with respect to the Poisson structure on $A(A_H^*S)$ induced from the Poisson structure on A_H^*S which is dual to the Lie algebroid $A_HS \rightarrow V$, and the Poisson structure on $A^2S \star AV$ dual to the Lie algebroid $A^2S \rightarrow AV$.*

PROOF. We will deduce these from the commutativity of Figures 9(a) and (b).




Figure 9.

In turn we will deduce these diagrams from the commutative diagram in Figure 10(a), which is valid for any manifold S [15, p.442]. Here $(\delta\nu)^\#$ is the map from forms to fields

induced by the canonical symplectic structure $\delta\nu$ on T^*S . The lower right triangle is the definition of Θ_S and the commutativity of the upper triangle is proved in [15, 9.6.7].

$$\begin{array}{ccc}
 T^*T^*S & \xrightarrow{R_{TS}} & T^*TS \\
 (\delta\nu)^\# \downarrow & \nearrow \Theta_S & \downarrow J_S^* \\
 TT^*S & \xrightarrow{I_{TS}} & T^\bullet TS
 \end{array}
 \quad (a)$$

$$\begin{array}{ccc}
 T^*T^*S & \xrightarrow{(\delta\nu)^\#} & TT^*S \\
 \downarrow & & \downarrow \\
 A_V^*T^*S & \xrightarrow{a_*} & TA_V^*S
 \end{array}
 \quad (b)$$

Figure 10.

For any $\mathcal{VB}$ -groupoid $(\Omega; G, A; M)$ there is a canonical isomorphism of double vector bundles $\mathfrak{R}_\Omega: A^*\Omega^* \rightarrow A^*\Omega$ which preserves the side bundles A and L^* and induces $-\text{id}$ on the cores A^*G [15, 11.5.13]. The maps $\tilde{R}_V$ and $\tilde{R}_H$ in Figure 9 are $\mathfrak{R}_\Omega$ for $\Omega = A_V^*S$ and $\Omega = A_H^*S$.

The maps Λ_H and Λ_V are the maps denoted $\mathcal{D}_H$ and $\mathcal{D}_V$ in [18, §3]. We recall the definition.

The cotangent groupoid of the groupoid $S \rightrightarrows H$ is $T^*S \rightrightarrows A_V^*S$. This is a symplectic groupoid and so the induced map $(\delta\nu)^\#: T^*T^*S \rightarrow TT^*S$ is an isomorphism of Lie groupoids as shown in Figure 10(b), where a_* denotes the anchor for the Lie algebroid structure on $A_V^*T^*S \rightarrow A_V^*S$.

$$\begin{array}{ccc}
 A_V^*T^*S & \xRightarrow{\quad} & A^*T^*K \\
 \downarrow & & \downarrow \\
 A_V^*S & \xRightarrow{\quad} & A^*K
 \end{array}
 \quad (a)$$

$$\begin{array}{ccc}
 TA_V^*S & \xRightarrow{\quad} & TA^*K \\
 \downarrow & & \downarrow \\
 A_V^*S & \xRightarrow{\quad} & A^*K
 \end{array}
 \quad (b)$$

$$\begin{array}{ccc}
 A_V^*TS & \xRightarrow{\quad} & A^*TK \\
 \downarrow & & \downarrow \\
 A_V^*S & \xRightarrow{\quad} & A^*K
 \end{array}
 \quad (c)$$

Figure 11.

This a_* is itself a morphism of double structures (indeed, of triple structures); the domain and target are shown in Figures 11(a) and (b). The $\mathcal{VB}$ -groupoid (a) arises by taking the vertical Lie algebroid of the double Lie groupoid $(T^*S; A_V^*S, A_H^*S; A^*K)$ and then dualizing; it has core $A^*(A_H^*S)$. The $\mathcal{VB}$ -groupoid (b) is the tangent of $A_V^*S \rightrightarrows A^*K$ and so has core $A(A_V^*S)$. We define Λ_H to be the core map of a_* . The definition of Λ_V is similar.

Both R_{TS} and Θ_S are also groupoid morphisms. For any $\mathcal{VB}$ -groupoid $(\Omega; G, A; M)$, R_Ω is a groupoid morphism over $\mathfrak{R}_\Omega: A^*\Omega^* \rightarrow A^*\Omega$ [15, 11.5.14]. In particular, R_{TS} is a Lie groupoid morphism over $\mathfrak{R}_{TS/TH}: A_V^*T^*S \rightarrow A_V^*TS$ where the subscript indicates that $\mathfrak{R}$ refers to the $\mathcal{VB}$ -groupoid $(TS; TH, S; H)$.

Similarly, Θ_S is a groupoid morphism over $\vartheta_{S/H}: TA_V^*S \rightarrow A_V^*TS$ [15, 11.5.11]. From the commutativity of Figure 10(a), it follows that

$$\vartheta_{S/H} \circ a_* = \mathfrak{R}_{TS/TH}.$$

It further follows that the core maps of $\vartheta_{S/H}$, a_* and $\mathfrak{R}_{TS/TH}$ commute. The core map of a_* is Λ_H by definition. To see the core of Figure 11(c), consider the triple structure in Figure 12(a). The core of the top face here is $A_V A_H S = A_2 S$ and the core of the bottom face is AH . Taking the dual of $A_V TS$ over TH produces the triple structure shown in Figure 12(b). The bundle $A_2 S \rightarrow AH$ of the top and bottom cores of (a) is parallel to the axis of dualization and so the bundle of top and bottom cores of (b) is $A_2 S \star AH \rightarrow AH$. The process of dualizing a triple vector bundle is described in detail in [23] and the modifications to take account of groupoid structures are straightforward.




Figure 12.

It is routine to verify that the core map of $\mathfrak{R}_{TS/TH}$ is $\tilde{R}_H$ and the core map of $\vartheta_{S/H}$ is $\tilde{\Theta}_H$. This proves the commutativity of the upper triangle in Figure 9(a). The proof for (b) is similar.

Now $R_{TS}: T^* T^* S \rightarrow T^* TS$ is an antisymplectomorphism and so $\mathfrak{R}_{TS/TH}$ is anti-Poisson [15, 11.5.16]. Likewise $(\delta\nu)^\#: T^* T^* S \rightarrow TT^* S$ is an antisymplectomorphism and so $\vartheta_{S/H}$ is anti-Poisson. It follows that the core maps are also anti-Poisson, and so Θ_H is a Poisson map. The case of Θ_V follows in the same way. This concludes the proof of Proposition 6.3 and with it the proof of Theorem 6.2. $\square$

It is also possible to prove that for any $\mathcal{L}\mathcal{A}$ -groupoid $(\Omega; G, A; M)$ applying the Lie functor yields a double Lie algebroid $(A\Omega; AG, A; M)$. In this case there is no version of the canonical involution J and it is necessary to work with prolonged Lie algebroid structures; see [20, 3.3].

Several examples of the double Lie algebroid of specific double Lie groupoids are given in [20, §4]. We conclude by noting three other examples.

Example 6.4 Take S to be the double groupoid $(M^4; M \times M, M \times M; M)$ in which a quadruple of points is regarded as defining a square by its four vertices. Then $A_2 S$ is the double tangent bundle $T^2 M$ and $\tilde{\Theta}_H$ reduces to Θ_M . The associated Lie bialgebroid is $(T^* T^* M, TT^* M)$ on base $T^* M$ with its standard symplectic structure.

Example 6.5 An *affinoid* W [37] on a manifold M may be regarded as a vacant double Lie subgroupoid of M^4 in the previous example, such that both side groupoids H and V are the graphs of simple foliations defined by surjective submersions $\pi_1: M \rightarrow Q_1$ and $\pi_2: M \rightarrow Q_2$

[16, §3]. The corresponding double Lie algebroid was calculated directly in [19] to be a double Lie subalgebroid $(A_2W; T^{\pi_1}M, T^{\pi_2}M; M)$ of T^2M with brackets defined by a pair of conjugate flat partial connections on M adapted to the two foliations. The Lie bialgebroid in this case is $(T^{\pi_1}M \ltimes (T^{\pi_2}M)^*, (T^{\pi_1}M)^* \rtimes T^{\pi_2}M)$ where the semi-direct structures are also defined by the connections.

Example 6.6 Let $(S; H, V; M)$ be a double Lie groupoid and write $D = A_2S$. Consider the cotangent double groupoid $(T^*S; A_V^*S, A_H^*S; A^*K)$. The double Lie algebroid of T^*S can be identified canonically with $(T^*D; D \star A(A_V^*S), D \star A(A_H^*S); A^*K)$, the cotangent double of D as shown in the top face of Figure 3(a).

References

- [1] A. L. Besse. *Manifolds all of whose Geodesics are Closed*, volume 93 of *Ergebnisse des Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1978.
- [2] V. Chari and A. Pressley. *A Guide to Quantum Groups*. Cambridge University Press, Cambridge, 1994.
- [3] T. J. Courant. Tangent Dirac structures. *J. Phys. A*, 23:5153–5160, 1990.
- [4] V. G. Drinfel’d. Quantum groups. In A. M. Gleason, editor, *Proceedings of the International Congress of Mathematicians, Berkeley, 1986*, pages 798–820. American Mathematical Society, Providence, R. I., 1987.
- [5] V. G. Drinfel’d. On Poisson homogeneous spaces of Poisson–Lie groups. *Theo. Math. Phys.*, 95(2):226–227, 1993.
- [6] C. Ehresmann. *Catégories et Structures*. Dunod, 1965.
- [7] J. Huebschmann. Differential Batalin–Vilkovisky algebras arising from twilled Lie–Rinehart algebras. In *Poisson geometry (Warsaw, 1998)*, pages 87–102. Polish Acad. Sci., Warsaw, 2000.
- [8] K. Konieczna and P. Urbański. Double vector bundles and duality. *Arch. Math. (Brno)*, 35(1):59–95, 1999.
- [9] Y. Kosmann–Schwarzbach. Exact Gerstenhaber algebras and Lie bialgebroids. *Acta Appl. Math.*, 41:153–165, 1995.
- [10] Y. Kosmann–Schwarzbach and F. Magri. Poisson–Lie groups and complete integrability. I. Drinfel’d bialgebras, dual extensions and their canonical representations. *Ann. Inst. H. Poincaré Phys. Théor.*, 49(4):433–460, 1988.
- [11] Z.–J. Liu, A. Weinstein, and Ping Xu. Manin triples for Lie bialgebroids. *J. Differential Geom.*, 45:547–574, 1997.
- [12] J.–H. Lu. Poisson homogeneous spaces and Lie algebroids associated to Poisson actions. *Duke Math. J.*, 86:261–304, 1997.

- [13] J.-H. Lu and A. Weinstein. Groupoïdes symplectiques doubles des groupes de Lie–Poisson. *C. R. Acad. Sci. Paris Sér. I Math.*, 309:951–954, 1989.
- [14] J.-H. Lu and A. Weinstein. Poisson Lie groups, dressing transformations, and Bruhat decompositions. *J. Differential Geom.*, 31:501–526, 1990.
- [15] K. Mackenzie. *General Theory of Lie Groupoids and Lie Algebroids*. London Mathematical Society Lecture Note Series, no. 213. Cambridge University Press, Cambridge, 2005.
- [16] K. C. H. Mackenzie. Double Lie algebroids and second-order geometry, I. *Adv. Math.*, 94(2):180–239, 1992.
- [17] K. C. H. Mackenzie. Drinfel'd doubles and Ehresmann doubles for Lie algebroids and Lie bialgebroids. *Electron. Res. Announc. Amer. Math. Soc.*, 4:74–87, 1998.
- [18] K. C. H. Mackenzie. On symplectic double groupoids and the duality of Poisson groupoids. *Internat. J. Math.*, 10:435–456, 1999.
- [19] K. C. H. Mackenzie. Affinoid structures and connections. In J. Grabowski and P. Urbański, editors, *Poisson Geometry: Stanisław Zakrzewski In Memoriam*, number 51 in Banach Center Publications, pages 175–186, Warsaw, 2000.
- [20] K. C. H. Mackenzie. Double Lie algebroids and second-order geometry, II. *Adv. Math.*, 154:46–75, 2000.
- [21] K. C. H. Mackenzie. Notions of double for Lie algebroids, . arXiv:math.DG/0011212.
- [22] K. C. H. Mackenzie. A unified approach to Poisson reduction. *Lett. Math. Phys.*, 53:215–232, 2000.
- [23] K. C. H. Mackenzie. Duality and triple structures. In *The breadth of symplectic and Poisson geometry*, volume 232 of *Progr. Math.*, pages 455–481. Birkhäuser Boston, Boston, MA, 2005.
- [24] K. C. H. Mackenzie and T. Mokri. Locally vacant double Lie groupoids and the integration of matched pairs of Lie groupoids. *Geom. Dedicata*, 77:317–330, 1999.
- [25] K. C. H. Mackenzie and Th. Th. Voronov. The double of a Lie bialgebroid is a double Lie bialgebroid. In preparation.
- [26] K. C. H. Mackenzie and Ping Xu. Lie bialgebroids and Poisson groupoids. *Duke Math. J.*, 73(2):415–452, 1994.
- [27] K. C. H. Mackenzie and Ping Xu. Integrability of Lie bialgebroids. *Topology*, 39:445–467, 2000.
- [28] S. Majid. Matched pairs of Lie groups associated to solutions of the Yang–Baxter equations. *Pacific J. Math.*, 141:311–332, 1990.
- [29] S. Majid. *Foundations of Quantum Group Theory*. Cambridge University Press, 1995.

- [30] R. A. Mehta. *Supergroupoids, double structures, and equivariant cohomology*, Thesis, Univ. California, Berkeley, 2006. [arXiv:math.DG/0605356](#)
- [31] T. Mokri. Matched pairs of Lie algebroids. *Glasgow Math. J.*, 39:167–181, 1997.
- [32] J. Pradines. Remarque sur le groupoïde cotangent de Weinstein–Dazord. *C. R. Acad. Sci. Paris Sér. I Math.*, 306:557–560, 1988.
- [33] D. Roytenberg. Courant algebroids, derived brackets and even symplectic supermanifolds. Thesis, Univ. California, Berkeley, 1999. [arXiv:math.DG/9910078](#).
- [34] D. Roytenberg and A. Weinstein. Courant algebroids and strongly homotopy Lie algebras. *Lett. Math. Phys.*, 46(1):81–93, 1998.
- [35] Th. Th. Voronov. Mackenzie theory and Q-manifolds. [arXiv:math.DG/0608111](#).
- [36] A. Weinstein. Coisotropic calculus and Poisson groupoids. *J. Math. Soc. Japan*, 40:705–727, 1988.
- [37] A. Weinstein. Affine Poisson structures. *Internat. J. Math.*, 1:343–360, 1990.